

Clinical Practice Guidelines for Oropharyngeal Dysphagia

A list of authors and their affiliations appears at the end of the paper

Objective: Dysphagia is a common clinical condition characterized by difficulty in swallowing. It is sub-classified into oropharyngeal dysphagia, which refers to problems in the mouth and pharynx, and esophageal dysphagia, which refers to problems in the esophageal body and esophagogastric junction. Dysphagia can have a significant negative impact on one's physical health and quality of life as its severity increases. Therefore, proper assessment and management of dysphagia are critical for improving swallowing function and preventing complications. Thus a guideline was developed to provide evidence-based recommendations for assessment and management in patients with dysphagia.

Methods: Nineteen key questions on dysphagia were developed. These questions dealt with various aspects of problems related to dysphagia, including assessment, management, and complications. A literature search for relevant articles was conducted using Pubmed, Embase, the Cochrane Library, and one domestic database of KoreaMed, until April 2021. The level of evidence and recommendation grade were established according to the Grading of Recommendation Assessment, Development and Evaluation methodology.

Results: Early screening and assessment of videofluoroscopic swallowing were recommended for assessing the presence of dysphagia. Therapeutic methods, such as tongue and pharyngeal muscle strengthening exercises and neuromuscular electrical stimulation with swallowing therapy, were effective in improving swallowing function and quality of life in patients with dysphagia. Nutritional intervention and an oral care program were also recommended.

Conclusion: This guideline presents recommendations for the assessment and management of patients with oropharyngeal dysphagia, including rehabilitative strategies. (JKDS 2023;13:77-106)

Keywords: Deglutition disorders, Rehabilitation, Diagnosis, Treatment, Therapeutics

INTRODUCTION

Dysphagia is a common clinical condition characterized by difficulty in swallowing. It is sub-classified into oropharyngeal dysphagia, which refers to problems in the mouth and pharynx, and esophageal dysphagia, which refers to problems in the esophageal

body and esophagogastric junction¹. Oropharyngeal dysphagia is characterized by difficulty in initiating a swallow or passing food through the mouth or throat², whereas esophageal dysphagia accompanies structural or inflammatory abnormalities or motility disorders³.

Dysphagia is associated with an acquired health condition, such as stroke, Parkinson's disease, or motor

Received: June 12 2023, Revised: June 22 2023, Accepted: July 7 2023, Published Online: July 30 2023

Corresponding author: Kyoung Hyo Choi, Department of Rehabilitation Medicine, Asan Medical Center, University of Ulsan College of Medicine, 88 Olympic-ro 43-gil, Seoul 05505, Korea

Tel: +82-2-3010-3800, Fax: +82-2-3010-6964, E-mail: kyounghyochoi@gmail.com

This clinical practice guideline was developed jointly by the Korean Academy of Rehabilitation Medicine and the Korean Dysphagia Society and was published in *Annals of Rehabilitation Medicine* and the *Journal of Korean Dysphagia Society* simultaneously on July 30, 2023.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Copyrights © The Korean Dysphagia Society, 2023.

neuron disease, as well as developmental disabilities. The prevalence of dysphagia is estimated to be 8% of the world's population, and its prevalence increases in the older adult population⁴. The prevalence of dysphagia in older people dwelling in communities is approximately 15% and approximately 30% in hospitalized patients¹. It occurs most commonly in old patients with neurological disorders and dementia, with a prevalence of 64% and 80%, respectively⁵.

Dysphagia can have a significant negative impact on one's physical health and quality of life as its severity increases⁶. If dysphagia is not properly evaluated and timely treated, it may worsen the quality of life and cause serious complications, such as dehydration, malnutrition, and aspiration pneumonia. Importantly, oropharyngeal dysphagia should be identified promptly considering the risk of aspiration. Various interventions such modifying food textures, positioning modification, or rehabilitative and compensatory strategies are designed to improve swallowing efficiency and reduce the risk of complications in patients with dysphagia⁷.

1. Purpose of clinical practice guidelines

The purpose of this clinical practice guideline (CPG) was to provide a guideline supported by scientific evidence for physicians and other healthcare professions who diagnose and treat patients with symptoms of dysphagia. This CPG is aimed to assist in decision-making for appropriate treatment options to improve the clinical outcome of patients with dysphagia and reduce extravagant costs to patients and the overall health care system.

2. Scope of CPGs

Assessment and management of dysphagia are mainly addressed in this CPG. Patients with progressive neurological disease and under 18 years of age were excluded from the scope of this CPG. This guideline does not aim to limit physicians' medical practices and is not used to evaluate the quality of their practices.

METHODS

1. CPG development group

The development group involved a development committee and an advisory committee, including three methodology experts. The development committee members consisted of 45 physicians (25 psychiatrists, 15 otorhinolaryngologists, 2 gastro-enterologists, and 3 dentists), 2 nursing staff, 3 nutritionists, 3 occupational therapists, and 2 language therapists.

For each key question (KQ), 55 development committee members determined the level of evidence and recommendation level.

2. KQs

This CPG consists of four categories: (1) assessment, (2) treatment, (3) nutritional management, and (4) complications and others. KQ was determined based on the Population Intervention Comparator Outcome (PICO) framework. According to the PICO strategy, adult patients of the population group who have symptoms or diagnosis of oropharyngeal dysphagia were included in this guideline. The number of KQs was determined by the number of interventions. The majority of outcomes focused mainly on improvement of dysphagia. If necessary, KQs were further divided into sub-KQs. Finally, a total of 19 KQs were formulated for this guideline.

3. Search strategy

A literature search was conducted for relevant articles using PubMed, Embase, the Cochrane Library, and one domestic database of KoreaMed, until April 2021. After establishing a highly sensitive strategy in combination with the natural language, the MeSH term was also used for PubMed and Cochrane Library, and the Emtree term was used for Embase. (Supplementary Data 1)

The search results were collated in EndNote. For each KQ, two independent reviewers excluded articles that did not meet the inclusion criteria of this CPG after reading titles and abstracts. Furthermore, full-

text assessments were followed to reject those that did not fulfill the inclusion criteria. Articles that included patients with progressive neurological diseases (e.g., Parkinson's disease or dementia) and patients who were under 18 years of age were excluded. We also included articles written in languages other than English or Korean, articles that exist only in abstract form, case reports, technical reports, and animal studies. Study screening and data extraction were independently performed by two reviewers. The reviewers attempted to resolve any disagreement by consensus. If necessary, the opinion of a third reviewer was put into consideration to resolve the disagreement.

Quality assessment was performed on selected articles. The bias assessment was conducted using Cochrane risk-of-bias 2.0 for randomized controlled trials (RCTs) and the risk-of-bias assessment tool for non-randomized studies for non-RCTs^{8,9}. The methodological quality of the systematic reviews (SRs) was evaluated using AMSTAR 1.0—a measurement tool to assess the methodological quality of SRs¹⁰. (Supplementary Data 2) The level of evidence and recommendation grade were established according to the Grading of Recommendation Assessment, Development and Evaluation (GRADE) methodology¹¹. The level of evidence was evaluated by assessing the degree of bias, consistency, directness, accuracy, and publication bias in the RCTs, non-RCTs, and SRs. The level of evidence for each KQ was based on the GRADE methodology as “high,” “medium,” “low,” and “very low.” (Table 1) The level of recommendation was determined according to a modified GRADE methodology divided into four levels. (Table 2) Several factors were considered, including the level of evidence, balance of benefits and harms, values and preferences, obstacles and facilitating factors, resource and cost, and clinical applicability. The KQs that could not be further developed due to poor existing research were represented as expert consensus.

For each KQ, at least two members of the development group participated in formulating and reviewing the draft recommendations. The working members continued discussions to reach consensus, and revisions

were made there after in accordance with the opinions of the advisory committee. The recommendations were also revised through a review process via e-mail and a wired meeting with experts in the relevant field, and a formal consensus was achieved. The level of evidence and recommendations for 19 KQs were evaluated, and the content of the recommendations and the recommendation grade was adjusted through in-depth discussion. The degree of consent for each committee member was selected from one (non-acceptance) to nine (acceptance) on a nine-point scale. If the score was ≥ 7 , consent was considered to be present. If at least 75% of the committee members agreed to the final version of the recommendations, it was deemed to have reached a consensus. Finally, the final version of the 19 recommendations was accepted. This guideline will be revised every 5 years, when there is solid evidence that it can affect the management and treatment of patients with oropharyngeal dysphagia.

RESULTS

1. Summary of recommendations

1) Assessment of oropharyngeal dysphagia

- KQ 1. Early screening
 - A. Evidence level: high
 - B. Grade of recommendation: strong
- KQ 2. Standardized screening test
 - A. Evidence level: not applicable
 - B. Grade of recommendation: expert consensus
- KQ 3. Videofluoroscopic swallowing study (VFSS) vs. clinical evaluation
 - A. Evidence level: moderate
 - B. Grade of recommendation: strong
- KQ 4. VFSS vs. fiberoptic endoscopic examination of swallowing (FEES)
 - A. Evidence level: inconclusive
 - B. Grade of recommendation: inconclusive

Table 1. Grading of Recommendations Assessment, Development and Evaluation quality level of evidence and meaning.

Quality level	Definition
High	We are confident that the estimate of the effect is close to the actual effect
Moderate	The estimates of the effect appear to be close to the actual effect but may vary considerably
Low	The confidence in the estimate of the effect is limited. The actual effect may differ significantly from the estimate of the effect
Very low	There is little confidence in the estimate of the effect. The actual effect will differ significantly from the estimate of the effect

Table 2. Grading of recommendations.

Strength of recommendations	Definition
Strong recommendation	The intervention/diagnostic test can be strongly recommended in most clinical practice, considering greater benefit than harm, evidence level, value and preference, and resources
Conditional recommendation	The intervention/diagnostic test can be conditionally recommended in clinical practice considering the balance of benefit and harm, evidence level, value and preference, and resources
Against recommendation	The harm caused by the intervention/diagnostic test maybe greater than its benefits. Moreover, considering the evidence level, value and preference, and resources, the intervention should not be recommended
Inconclusive	It is not possible to determine the strength and direction of recommendation because of a very low or insufficient evidence level, uncertain or variable balance of benefit and harm, value and preference, and resources
Expert consensus ^{a)}	Although clinical evidence is insufficient, it is recommended to be used in accordance with clinical experience and expert consensus when considering the benefits and risks of the treatment, the level of evidence, values and preferences, and resources

Each statement is shown as a combination of the strength of recommendations and level of evidence.

^{a)}In the case of a consensus statement by an expert opinion, the recommendation grade and level of evidence are not indicated.

2) Treatment for oropharyngeal dysphagia

KQ 5. Oropharyngeal sensory stimulation

A. Evidence level: low

B. Grade of recommendation: conditional

KQ 6. Exercises

KQ 6.1. Tongue and pharyngeal muscle strengthening exercise

A. Evidence level: moderate

B. Grade of recommendation: strong

KQ 6.2. Expiratory muscle strength training (EMST)

A. Evidence level: low

B. Grade of recommendation: conditional

KQ 7. Compensatory swallowing technique

A. Evidence level: conditional

B. Grade of recommendation: very low

KQ 8. Neuromuscular electrical stimulation (NMES)

KQ 8.1. Swallowing therapy with or without NMES in non-progressive neurological disease

A. Evidence level: moderate

B. Grade of recommendation: strong

KQ 8.2. Swallowing therapy with or without NMES in head and neck cancer

A. Evidence level: low

B. Grade of recommendation: conditional

KQ 8.3. Pharyngeal electrical stimulation (PES) in non-progressive neurological disease

A. Evidence level: low

B. Grade of recommendation: conditional against

KQ 9. Stimulation of the transient receptor potential (TRP) channel with drugs

A. Evidence level: moderate

B. Grade of recommendation: conditional

- KQ 10. Biofeedback training
 A. Evidence level: low
 B. Grade of recommendation: conditional
- KQ 11. Specific treatment for cricopharyngeal (CP) dysfunction
 KQ 11.1. CP botulinum toxin injection
 A. Evidence level: low
 B. Grade of recommendation: conditional
- KQ 11.2. CP myotomy
 A. Evidence level: not applicable
 B. Grade of recommendation: expert consensus
- KQ 11.3. Balloon dilatation
 A. Evidence level: moderate
 B. Grade of recommendation: conditional
- KQ 12. Swallowing education
 A. Evidence level: very low
 B. Grade of recommendation: conditional
- KQ 13. Noninvasive brain stimulation
 KQ 13.1. Transcranial direct current electrical stimulation (tDCS)
 A. Evidence level: low
 B. Grade of recommendation: conditional
- KQ 13.2. Repetitive transcranial magnetic stimulation (rTMS)
 A. Evidence level: very low
 B. Grade of recommendation: conditional

3) Nutrition for oropharyngeal dysphagia

- KQ 14. Tube feeding in patients with suspected long-term dysphagia
 A. Evidence level: very low
 B. Grade of recommendation: conditional
- KQ 15. Modifying food textures
 A. Evidence level: very low
 B. Grade of recommendation: conditional
- KQ 16. Nutritional intervention
 A. Evidence level: moderate
 B. Grade of recommendation: strong

4) Complications and others

- KQ 17. Incidence and mortality rates of aspiration pneumonia
 A. Evidence level: high
 B. Grade of recommendation: strong

- KQ 18. The effect of oral care program
 A. Evidence level: moderate
 B. Grade of recommendation: strong
- KQ 19. The effect of multidisciplinary team approach
 A. Evidence level: low
 B. Grade of recommendation: conditional

2. Assessment of oropharyngeal dysphagia

KQ 1. Is early screening effective in improving the prognosis in patients with suspected oropharyngeal dysphagia?

Dysphagia is a medical condition that increases the risk of various complications, such as dehydration, malnutrition, aspiration pneumonia, and airway obstruction, which can lead to serious disability or even death¹²⁻¹⁴. Various methods have been used to screen patients with dysphagia, and it is important to investigate whether early screening tools for dysphagia are effective in improving prognosis and preventing complications that can occur due to dysphagia.

One RCT conducted by Schmidt Leuenberger et al.¹² reported that the incidence of pneumonia decreased in patients who received a clinical assessment of dysphagia (early screening) after pulmonary resection. Ten retrospective studies, including patients with stroke, post extubation dysphagia in an intensive care unit (ICU), and traumatic cervical injuries, showed that respiratory complications were significantly reduced after the early screening of dysphagia¹²⁻²¹. Considering that the above studies showed consistent results, early screening is recommended in patients with suspected oropharyngeal dysphagia to reduce the occurrence of pneumonia. If patients with oropharyngeal dysphagia are screened early and are provided with appropriate treatment as early as possible, serious complications, such as pneumonia, can be prevented.

Recommendation

Early screening is strongly recommended in patients with suspected oropharyngeal dysphagia to

reduce the occurrence of pneumonia with high levels of evidence.

KQ 2. Is the standardized screening test more effective for the diagnosis of dysphagia than a single screening test?

If screening and appropriate treatments for dysphagia are provided as early as possible, the occurrence of complications related to the disease can be reduced²². Various screening tests for dysphagia have been developed and used in clinical practice. Single screening tests, such as the 3-oz water swallow test or the volume-viscosity swallow test, are used, where food is swallowed directly to check the presence of aspiration by coughing, voice change, and change in oxygen saturation²³. Standardized screening tests, such as the Burke dysphagia screening test (BDST), Gugging Swallowing Screening Test (GUSS), Standardized Swallowing Assessment (SSA), Toronto Bedside Swallowing Screening Test (TOR-BSST), and Clinical Functional Scale for Dysphagia, use a clinical scale by combining various clinical items²⁴⁻²⁶.

Shin et al.²³ compared the screening abilities of the single screening test (3-oz water test) and standardized screening tools (GUSS, BDST, and SSA) based on the VFSS findings in patients with stroke. The results showed that there was no significant difference in the screening ability of GUSS compared to other screening tests. Lopes et al.²⁷ also compared the water test and GUSS and reported that there were no differences in the occurrence of stroke-associated pneumonia, mortality rate, ICU admission rate, and functional status between the two groups. Both studies reported that there was no difference in the accuracy of diagnosis or the occurrence of complications between standardized and single screening tests. However, a standardized screening test tool can evaluate a patient's swallowing function without directly swallowing food. Therefore, for high-risk patients who cannot swallow their saliva properly, a standardized screening test seems safer than a single screening test that forces them to swallow water.

Recommendation

A standardized screening test may be considered to diagnose dysphagia in patients with suspected oropharyngeal dysphagia (expert consensus).

KQ 3. Is VFSS more effective than clinical evaluation in diagnosing oropharyngeal dysphagia?

Oropharyngeal dysphagia can be evaluated using various clinical assessment tools that can be performed at the bedside, as well as more comprehensive diagnostic tests, such as VFSS, can be applied. VFSS is considered a gold-standard evaluation tool for dysphagia, because it can visualize a series of swallowing processes occurring in the oral, pharyngeal, and esophageal phases²⁸.

If VFSS is performed in patients with oropharyngeal dysphagia, the presence of dysphagia can be detected more accurately than during a clinical evaluation. Two SRs^{29,30} and four studies³¹⁻³⁴ that compared the effectiveness of clinical evaluation and VFSS were identified. Both SRs that included patients with stroke showed that no statistically significant differences were found between clinical evaluation and VFSS in predicting the occurrence of aspiration pneumonia^{29,30}. Other studies recommended that VFSS is cost-effective and superior when compared to a clinical bedside swallowing evaluation³¹⁻³⁴. Therefore, clinical evaluation can be useful as an early screening test, and VFSS can be additionally performed to accurately diagnose the presence of dysphagia.

Recommendation

VFSS is strongly recommended for diagnosis of dysphagia with moderate levels of evidence.

KQ 4. Is FEES more effective than VFSS for the diagnosis of dysphagia?

It is necessary to diagnose patients with oropharyngeal dysphagia accurately in order to prevent detrimental complications, such as aspiration pneumonia, and to provide sufficient nutrition as early as possible. In addition, a diagnostic test for dysphagia

is necessary to evaluate the recovery of swallowing function before and after treatment.

Diagnostic tests for dysphagia include bedside screening, VFSS, and FEES. VFSS is widely used as a standard diagnostic tool for oropharyngeal dysphagia, and FEES is applied to visualize the swallowing function; it can easily be performed repeatedly, even while lying down. Recently, FEES has been frequently performed to diagnose oropharyngeal dysphagia, and many studies on its usefulness have been reported.

One RCT conducted by Aviv³⁵ compared the effectiveness of VFSS and FEES in 126 patients with dysphagia. The results showed that there were no statistically significant differences in determining the prevalence of aspiration pneumonia between patients who received FEES and VFSS. A study conducted by Wu et al.³⁶ reported that FEES was more sensitive in detecting penetration, aspiration, pharyngeal retention, and cough reflex, while Fattori et al.³⁷ recommended that FEES was useful in visualizing pharyngeal residue. FEES showed superior results in observing airway penetration and pharyngeal residue than VFSS^{38,39}.

Similar levels of safety and efficacy have been observed between FEES and VFSS in patients with dysphagia. There is no radiation exposure, and it can be easily inspected even in medical institutions that do not have fluoroscopy equipment. The endoscopic swallowing test has a high diagnostic benefit since laryngeal movement can be detected during actual swallowing when dysphagia is suspected. In addition, it has the advantage of being performed repeatedly in various places, and there are no significant complications nor patient discomfort during the procedure. However, it is difficult to observe the whole swallowing process using FEES. The phases of swallowing cannot be assessed, especially the oral phase.

Recommendation

There is insufficient evidence to conclude whether FEES is more effective than VFSS. The two test methods can be performed complementary to each other.

KQ 5. Is oropharyngeal sensory stimulation therapy effective in improving swallowing function and quality of life?

Sensory stimulation therapy is believed to be a potential strategy for treating dysphagia as it activates the peripheral sensory nerves in the larynx and the pharynx to protect the airway from aspiration. It has been reported that the use of NMES can improve swallowing in patients with dysphagia by stimulating the afferent nerves and increasing the sensory input to the central nervous system⁴⁰.

A RCT conducted by Maeda et al.⁴¹ showed that 20 patients with oropharyngeal dysphagia who received sensory stimulation showed improvement in oral nutritional intake and functional oral intake scale (FOIS). Zhang et al.⁴² also conducted a RCT and compared the effectiveness of the traditional swallowing therapy, sensory approach (NMES on the sensory input) combined with traditional swallowing therapy, and motor approach (NMES on the motor muscle) combined with traditional swallowing therapy. All the groups showed improvement in swallowing function and quality of life after 4 weeks of treatment, but the sensory approach combined with traditional swallowing therapy showed a statistically significant improvement compared to the other groups⁴². Another RCT performed by Rofes et al.⁴³ showed that after 10 days of treatment with sensory or motor surface electrical stimulation, 20 patients with chronic post stroke dysphagic (10 sensory vs. 10 motor) showed improvement in swallowing function after treatment compared to that before treatment. These studies suggest that oropharyngeal sensory stimulation therapy in patients with oropharyngeal dysphagia may contribute to an improvement in swallowing function and an increase in quality of life.

Recommendation

Oropharyngeal sensory stimulation therapy is recommended in patients with oropharyngeal dysphagia to improve swallowing function and improve quality of life.

KQ 6.1. Are tongue and pharyngeal muscle strengthening exercises effective in improving swallowing function and quality of life, reducing the incidence of pneumonia, and improving quality of life?

Tongue and pharyngeal muscle strengthening exercises are one of the treatment methods for dysphagia. Strengthening exercises are often applied for better lingual and pharyngeal strength and improvement in swallowing ability⁴⁴. These exercises aimed to increase the diameter of the upper esophageal sphincter (UES) opening and decrease post deglutitive aspiration and dysphagic symptoms⁴⁵.

Twelve RCTs investigated whether strengthening exercises for the tongue and laryngeal muscles were effective in improving the swallowing function and reducing the incidence of aspiration pneumonia compared to the control group. Three and eight RCTs on patients with brain lesions and head and neck cancer, respectively, were found. These studies applied rehabilitative techniques, such as head lift exercise, Shaker exercise, Mendelsohn maneuver, and muscle strengthening exercises of the tongue and larynx.

Regarding patients with stroke, in a RCT conducted by Kang and Kim¹, patients with dysphagia showed a significant improvement in the neck strength and swallowing function when head lift exercises were performed for 30 minutes, 5 times a week for 6 weeks. RCTs conducted by McCullough et al.⁴⁶ and McCullough and Kim⁴⁷ also showed that the 2 weeks of the Mendelsohn method training improved hyoid anterior and superior movements and increased UES opening and swallow physiology in 18 patients with stroke and dysphagia. Regarding patients with oropharyngeal cancer, Kotz et al.⁴⁸ investigated the effect of prophylactic swallowing exercises in 26 patients with head and neck cancer. This study showed that patients who performed swallowing exercises (five exercises, including effortful swallowing, super-supraglottic swallowing, tongue hold, tongue retraction, and Mendelsohn maneuver, 10 times per day, and three times a day) showed better swallowing outcomes than patients who did not perform the exercises at

3 and 6 months after the treatment⁴⁸. A RCT conducted by Lazarus et al.⁴⁴ also showed that tongue-strengthening exercises improved swallowing function in 12 patients with oropharyngeal cancer who underwent radiotherapy compared to 11 patients who were administered only conventional treatment. Notably, other studies also reported similar results.

In summary, tongue and pharyngeal muscle strengthening exercises contribute to improving swallowing function and reducing aspiration pneumonia in patients with brain lesions and head and neck tumors. Since these exercises can be easily performed for inpatients and outpatients in rehabilitation clinics, they are recommended for the treatment of dysphagia.

Recommendation

Tongue and pharyngeal muscle strengthening exercises are recommended to improve swallowing function and reduce the incidence of pneumonia.

KQ 6.2. Is EMST effective in improving swallowing function or quality of life?

The EMST increases subglottic air pressure, while changes in motor unit recruitment and neuromodulation have been proposed as the mechanism of effect.

Four RCTs on EMST showed that it improved swallowing function in patients with dysphagia. Three RCTs on patients with stroke⁴⁹⁻⁵¹ and one RCT on patients with multiple sclerosis⁵² used commercially available EMST 150 equipment (Aspire Products LLC., Cape Carteret, NC, USA) and compared the effect with the sham treatment. As for EMST, a protocol of five sets per day was applied for 5 days a week. One set included five vigorous exhalations at 70% of the maximum expiratory pressure. All of these studies showed that EMST was effective in improving swallowing function compared to the control group⁴⁹⁻⁵², and the study conducted by Silverman et al.⁵² also reported that it improved swallowing-related quality of life. EMST can improve swallowing function as well as respiratory function in patients with non-progressive neurological disease, thereby contributing to improving

swallowing-related quality of life, but it can be only applied to patients who can fully understand the training method. Clinicians should be aware of its side effects, as performing strong expiratory pressure can cause hypotension, hyperventilation, or dizziness.

Recommendation

EMSTs are suggested to improve swallowing function and quality of life.

KQ 7. Are compensatory swallowing maneuvers effective in improving swallowing function, lowering the incidence of pneumonia, and improving quality of life?

Compensatory maneuvers are one of the rehabilitative strategies for dysphagia management. The compensatory swallowing maneuvers aim to keep patients safe when swallowing and promote temporary stability without permanently changing the swallowing mechanism. Compensatory swallowing maneuvers include chin tuck, chin down, head extension, head rotation, effortful swallow, Mendelsohn maneuver, tongue-hold maneuver, or Masako maneuver, supraglottic swallow, and super-supraglottic swallow. Compensatory swallowing maneuvers can induce immediate stability of the swallowing; however, as the same method must be applied every time swallowing occurs, patients may get tired easily due to the repeated application.

According to the observational study conducted by Solazzo et al.⁵³, compensatory swallowing techniques, such as a chindown posture, head turned posture, and a hyperextended head posture, promoted safe swallowing in 66 (88.0%) of 75 patients with dysphagia. Furthermore, a study published by Ra et al.⁵⁴ also showed the effect of chin tuck, and the study conducted by Miyamoto et al.⁵⁵ reported that the chin-down maneuver was beneficial to swallowing function. The evidence for the effect of effortful swallowing seems to be weak at present, and no studies were found regarding the effect of Mendelsohn maneuver and tongue hold maneuver as compensatory swallowing techniques. Regarding supraglottic and super-supraglottic swallowing, a study

reported that super-supraglottic swallow changes the airway closure and hyoid-larynx movement⁵⁶, while another study reported that supraglottic swallow does not change the propelling pressure of food lumps⁵⁷. Further studies on their direct effect on swallowing function are needed in the future.

Compensatory swallowing maneuvers do not require a large burden of time and cost and can positively affect swallowing function and prevent complications. To date, there has been no RCT studying compensatory swallowing maneuvers; therefore, the level of evidence was considered very low.

Recommendation

The compensatory swallowing technique is suggested to improve the swallowing function.

KQ 8.1. Is surface NMES combined with swallowing therapy better for improving dysphagia in patients with non-progressive neurological disease compared with swallowing therapy alone?

Conventional swallowing therapy includes strengthening exercises for muscles involved in swallowing, compensatory maneuvers, sensory and tactile stimulation, and biofeedback⁵⁸. In 2001, U.S. Food and Drug Administration approved the use of NMES, such as VitalStim[®] (Chattanooga Group, Hixson, TN, USA), for dysphagia treatment. This treatment method involves attaching electrodes to the skin and transmitting electrical signals to the muscles involved in swallowing to induce contraction.

Various studies evaluated the combined effect of surface NMES, including 10 RCTs^{42,59-67} and two non-RCTs^{68,69}. In most studies, the combination of NMES and swallowing therapy significantly improved objective and subjective indicators of swallowing function. A RCT conducted by Lee et al.⁶² showed that when combined with NMES and conventional swallowing therapy, the FOIS was higher at 3 and 6 weeks after treatment compared to conventional swallowing therapy alone. A RCT conducted by Terré and Mearin⁶⁶ reported that the combination of NMES and conventional swallowing therapy improved oral intake

function and reduced aspiration in patients with dysphagia. A RCT conducted by Xia et al.⁶⁷ also showed that the combination of NMES and conventional swallowing therapy was conducive to recovery from poststroke dysphagia. Nine out of ten studies showed similar results, reporting that the combination of NMES and conventional swallowing therapy helped improve the swallowing questionnaire scores or swallowing test results more significantly than swallowing therapy alone. This treatment method is considered safe and effective. Therefore, NMES, in conjunction with swallowing therapy, is recommended for the treatment of dysphagia.

Recommendation

The combination of surface NMES and swallowing therapy is recommended for improving the swallowing function in patients with non-progressive neuropathic lesions.

KQ 8.2. Is surface NMES combined with swallowing therapy better for improving dysphagia compared with swallowing therapy alone in patients with head and neck cancer?

Dysphagia after head and neck cancer surgery depends on the type and extent of the damage to the structures removed during surgery. It can also occur after radiation therapy, which induces progressive fibrosis of muscles or soft tissues that leads to progressive dysphagia⁷⁰.

For the treatment of dysphagia after head and neck cancer surgery, strengthening exercises of remaining muscles, postural maneuvers, and biofeedback techniques have been used to replace the original functions of the resected structures and minimize the weakening of the muscles⁷⁰. The surface NMES has also been applied as an alternative treatment to strengthen the muscles through direct muscle contractions and prevent the atrophy of denervated muscles.

One RCT and two case-control studies investigated the effectiveness of the combination of NMES and conventional swallowing therapy. A RCT conducted by Ryu et al.⁷¹ showed that NMES combined with

traditional swallowing training (14 patients) was superior to traditional swallowing training alone (12 patients) in patients with head and neck cancer. Two case-control studies showed that NMES induced significant scores in the FOIS assessment and the degree of movement in the speed of the hyoid bone^{72,73}. The combination of NMES and swallowing therapy is beneficial for improving dysphagia in patients with head and neck cancer.

Recommendation

The combination of surface NMES and swallowing therapy is suggested for improving the swallowing function in patients with head and neck cancer.

KQ 8.3. Does PES improve swallowing function and prevent pneumonia in patients with dysphagia caused by non-progressive neurological disease?

PES, which provides electrical stimulation directly to the pharynx, induces activation of the pharyngeal motor cortex via the corticobulbar pathway⁷⁴. PES aims for cortical plasticity by activating the cerebral motor cortex by transmitting repetitive electrical stimulation into the pharynx.

Five RCTs on the PES compared its effect with sham stimulation⁷⁴⁻⁷⁸. Four RCTs reported that PES did not significantly improve the swallowing function or prevent pneumonia^{74,75,77,78}, while only Jayasekaran et al.⁷⁶ reported that PES was safe, reduced aspiration, and improved feeding status. Therefore, evidence of the positive effect of PES on improving swallowing function and preventing pneumonia is lacking.

Recommendation

It is difficult to recommend the application of PES in patients with oropharyngeal dysphagia caused by non-progressive nerve lesions since the effects of improving the swallowing function and preventing pneumonia are not clear.

KQ 9. Is stimulating the TRP channel with drugs effective in improving swallowing function, lowering the incidence of pneumonia, and improving the quality of life in patients with oropharyngeal dysphagia?

Although drug treatment for dysphagia is not currently widely applied in clinical practice, a number of drugs have been studied for their effect of improving swallowing response and preventing airway aspiration. TRP channel is a receptor that converts temperature stimuli into electrical signals and transmits them to the central nervous system. A number of substances, including a TRPV1 agonist (capsaicin) and a TRPM8 agonist (menthol), were used in several studies to induce swallowing responses in patients with dysphagia.

The effect of a capsaicin tablet was compared with a placebo tablet for 4 weeks before meals, and upper respiratory protective reflexes significantly improved in 64 older residents in a nursing home⁷⁹. When TRPV1 agonist (capsaicin 1×10^{-5} M) was administered for 10 days, the penetration-aspiration scale (PAS) score on the VFSS significantly decreased from 5.23 ± 2.04 to 3 ± 1.47 ($P=0.002$)^{80,81}. The stimulation of TRPV1 improved swallowing safety and shortened the swallow response in older adult patients with oropharyngeal dysphagia. In two RTCs conducted by Cui et al.⁸² and Wang et al.⁸³, capsaicin in conjunction with ice or thermal tactile stimulation was beneficial to the recovery of the swallowing function in patients with stroke and dysphagia. In addition to capsaicin, the effect of menthol (TRPM8 agonist) and piperine (TRPV1/TRPA1 agonist) injection studies on the swallowing function have also been reported^{84,85}. Most RCTs (11 out of 12) reported that TRP channel stimulation improved swallowing function in the short term, and four non-RCTs reported that TRP stimulation improved the swallow response and swallowing function.

Currently, little is known about pharmacological approaches for the treatment of oropharyngeal dysphagia. Stimulation of the TRP channel via TRPV agonists (including capsaicin), administered through the oral cavity, oropharynx, or through stimulation of the ear canal, showed improvement in the swallowing

function for a short period. However, its long-term use should be further investigated. The administration of a TRPV agonist, such as capsaicin, showed a short-term improvement in the swallowing effect, and it can be easily applied in Korea as capsaicin-rich red pepper is a very common food in Korea. The regular stimulation of the TRP channel seems to promote the recovery of the swallowing function in patients with dysphagia, but a clear standard for the dosage and the long-term use of drugs are needed in the future.

Recommendation

Pharmacological stimulation of the TRP channel is suggested for improving swallowing function.

KQ 10. Is biofeedback training effective for improving swallowing function, lowering the incidence of pneumonia, and improving quality of life?

Biofeedback is a technique that provides biological information to patients in real-time and facilitates normal movement patterns or induces sufficient muscle contractions during muscle-strengthening training⁸⁶. Using biofeedback, the exercise performance is notified to the patients through audio-visual information to empower effective rehabilitation. Neuromuscular biofeedback methods include electromyography (EMG) biofeedback and real-time ultrasound imaging biofeedback. EMG biofeedback measures myoelectric signals through surface electrodes attached to target muscles and converts them into visual and auditory signals to induce sufficient muscle contractions⁸⁷.

A RCT conducted by Shin et al.⁸⁸ showed that suprahyoid muscle activity improved after 4 weeks of biofeedback training using surface EMG, compared to that before treatment in 45 patients with dysphagia after stroke. The effect of visuoauditory biofeedback is superior to visual biofeedback or self-exercise alone⁸⁸. Another RCT conducted by Moon et al.⁸⁹ showed that swallowing training (effortful swallow and Mendelsohn maneuver) with surface EMG bio-

feedback training was more effective than swallowing training alone. Other case-control studies also showed that surface EMG biofeedback induced improvement in swallowing scores, such as functional dysphagia scale (FDS), PAS, videofluoroscopic dysphagia scale (VDS), or FOIS⁹⁰⁻⁹⁴.

In summary, swallowing treatment using biofeedback is an effective adjunct to conventional swallowing therapy to improve swallowing function in patients with dysphagia. Although there is not much evidence, the benefit is clear compared to the harm. Since there are many hospitals that do not have biofeedback equipment, if applicable, biofeedback in conjunction with conventional swallowing therapy is recommended.

Recommendation

Biofeedback training is suggested for improving swallowing function.

KQ 11.1. Is CP botulinum toxin injection effective for improving swallowing function, reducing the risk of aspiration pneumonia, and improving diet in patients with CP dysfunction?

The CP muscle, which is located between the throat and esophagus, is the main component of UES. CP dysfunction or impaired relaxation due to various diseases, such as neurological disease, head and neck cancer, and Zenker's diverticulum, causes symptoms of dysphagia, aspiration, and weight loss^{95,96}. In patients with CP dysfunction, the cricopharyngeus fails to open during swallowing. Treatments for CP dysfunction include swallowing therapy and interventional techniques, such as botulinum toxin injection, dilatation, and myotomy. The effectiveness of CP botulinum toxin injection, which was first introduced by Blitzer in 1993, has been reported in previous studies⁹⁷.

Several studies reported the effects of CP botulinum injection, including two studies conducted by Alfonsi et al.⁹⁸ and Kelly et al.⁹⁹, which included 69 and 49 patients with CP dysfunction, respectively. Scores obtained from the eating assessment tool and Dysphagia Outcome and Severity Scale (DOSS) showed

improvement after botulinum toxin injection. Botulinum toxin injection was highly effective in the treatment of dysphagia. In the study published by Jeong et al.¹⁰⁰ and Kim et al.¹⁰¹, the success rate of botulinum toxin injection was 63.9% and 78.6%, respectively. The complication rate was very low, with only one patient showing temporary unilateral vocal fold paralysis¹⁰¹. The relative risk of the CP botulinum toxin injection seems to be low, and the benefits seem greater than the harms.

Recommendation

CP botulinum toxin injection is suggested to be performed while carefully considering the characteristics of dysphagia of each patient and the advantages and disadvantages of treatment.

KQ 11.2. Is CP myotomy effective in improving swallowing function, reducing the risk of aspiration pneumonia, and improving diet in patients with CP dysfunction?

CP myotomy is one of the surgical interventions applied for the treatment of CP dysfunction. CP myotomy is a treatment method that induces functional improvement by performing a complete incision of the CP muscle. Two approaches have been developed: the external transcervical approach, which has been performed for a long time, and the endoscopic CP myotomy, which has been performed relatively recently. In patients with defective CP relaxation, adequate oral and pharyngeal propulsion and laryngeal elevation may have a positive effect on improving the swallowing function. However, it has been reported that the effect of CP myotomy is inconclusive for patients who complain of subjective symptoms without clear abnormalities in anatomy and function¹⁰².

McKenna and Dedo¹⁰³ conducted a study on the effects of CP myotomy through an external transcervical approach on 47 patients with CP dysfunction in 1992. After surgery, 45% and 30% of patients showed normal and improved swallowing function, respectively, in the clinical symptom evaluation. Brigand et al.¹⁰⁴ showed that among 253 patients with severe dysfun-

ctional pharyngo-esophageal junction who underwent CP myotomy through an external transcervical approach, 75% of patients reported improvement in oropharyngeal. In addition, postoperative complications were observed in 15.8% of the patients, with the most common complications being mucosal break and pulmonary infection. Dauer et al.¹⁰⁵ compared eight patients who underwent CP myotomy through an external transcervical approach and 14 patients who underwent endoscopic CP myotomy. The laser technique was as effective as the transcervical approach, with a low risk of major complications. Takes et al.¹⁰⁶ and Ho et al.¹⁰⁷ showed that most patients reported improvement in dysphagia symptoms after endoscopic CP myotomy at 3 and 6 months, respectively.

Symptomatic improvement after CP myotomy seems to be approximately 60%–70% in patients with CP dysfunction. However, the clinician should consider that CP myotomy is invasive, irreversible, and permanent treatment with a possible risk of complications. Therefore, it seems difficult to judge whether the risks will be significantly higher than the relative risk for benefits, so an individual approach based on the patient's condition seems necessary.

Recommendation

CP myotomy is suggested to be performed carefully in selective cases who receive refractory to conventional treatment while considering the potential side effects, advantages, and disadvantages of the treatment. An individual approach based on the patient's condition seems necessary.

KQ 11.3. Is balloon dilatation effective in improving swallowing function, reducing the risk of aspiration pneumonia, and improving diet in patients with CP dysfunction?

Balloon dilatation is a treatment method for CP dysfunction and is known as a relatively safe and efficacious in relieving UES dysfunction¹⁰⁸. One of the dilatation techniques is the use of bougies, which reduces UES pressure and increases relaxation.

Balloon dilatation for CP dysfunction restored UES resting pressure, improved UES relaxation, strengthened pharyngeal propulsion, and improved functional oral intake¹⁰⁸.

A RCT conducted by Wei et al.¹⁰⁹ showed that conventional swallowing therapy with modified balloon dilatation increased the excitability of affected projection and induced better improvement in FOIS compared to conventional swallowing alone at 3 weeks of treatment. When balloon dilatation was compared with laser myotomy, both improved UES opening for at least 6 months after the treatment¹¹⁰. Other studies also reported the effectiveness of balloon dilatation for treating CP dysfunction.

Balloon dilatation is a procedure that secures the visual field through an endoscope, and the risk of side effects is not high compared to other interventional procedures. It is considered valuable for treating CP dysfunction, but it is still invasive to a degree and may, therefore, require repeated procedures when CP dysfunction recurs.

Recommendation

Balloon dilatation is suggested to be performed carefully in selective cases that are refractory to conventional treatment, considering the advantages and disadvantages of the treatment. An individual approach based on the patient's condition seems necessary.

KQ 12. Is the swallowing education program effective for improving swallowing function, reducing the risk of aspiration pneumonia, and improving diet?

For the treatment of dysphagia, various methods, including electrical stimulation therapy and swallowing exercises, are effectively used in clinical practice for the recovery of swallowing function. Recently, the need for a comprehensive swallowing education program that includes an understanding of dysphagia, self-swallowing exercise, and management has been emphasized.

Kang et al.¹¹¹ applied a comprehensive bedside

swallowing exercise education program for 2 months in addition to the conventional swallowing therapy for patients with stroke and dysphagia. A video recording of the swallowing exercises, including oral, pharyngeal, laryngeal, and respiratory exercises, were played in the ward. These videos led to an improvement in swallowing function, emotional state, and quality of life in the patient group who received swallowing education. Cho et al.¹¹² also showed that the supervised self-exercise swallowing training program for 4 weeks induced improvement in the VDS in patients with stroke. Mashhour et al.¹¹³ showed that swallowing exercise programs are also effective in patients with dysphagia due to the presence of head and neck tumors during radiation therapy. In addition, Chen et al.¹¹⁴ confirmed that the swallowing exercise education program was effective in improving emotional dysphagia quality of life 6 months after treatment of oral cavity cancer.

The swallowing education program is effective in improving swallowing function. There is no harm that can be caused by performing the program, and the motivation for participation in the program is necessary to maximize its effect. The swallowing education program can be used as a modality for rehabilitation for patients with dysphagia. are mild and include local erythema, tingling, or itching. In conclusion, tDCS applied to the contralateral or bilateral hemispheres is beneficial for the improvement of swallowing function when combined with swallowing therapy in patients with dysphagia.

Recommendation

The comprehensive swallowing education program, including self-exercise swallowing training, is suggested for improving swallowing function.

KQ 13.1. Is tDCS effective in improving swallowing function, reducing the risk of aspiration pneumonia, and improving diet?

tDCS, one of the non-invasive brain stimulation treatment methods, aims to induce functional improvement through neuroplasticity by controlling the

activity of neural networks in the cortical areas. Positive therapeutic effects of tDCS on the recovery of language, motor, and cognitive function in patients with stroke have been reported in previous studies¹¹⁵. Regarding the effect of tDCS on patients with dysphagia, Jefferson et al.¹¹⁶ first investigated the applicability of anodal tDCS stimulation to the pharyngeal motor cortex, and later, several studies reported the effects of tDCS stimulation on dysphagia after stroke.

In 2011, Kumar et al.¹¹⁷ conducted a RCT wherein anodal tDCS was applied to the pharyngeal motor cortex area contralateral to the lesion site in patients with dysphagia with subacute stroke. There was a significant difference after anodal tDCS (seven patients) compared with sham tDCS (seven patients) in DOSS scores. Other RCTs also reported that anodal tDCS induced better swallowing function when comparing the effect of anodal tDCS with sham tDCS¹¹⁸⁻¹²². Most recently, in a RCT conducted by Sawan et al.¹²³, 20 patients who received anodal tDCS with conventional rehabilitation therapy for 5 days showed significant improvement in the swallowing function (VFSS and DOSS score) compared with 20 patients who received sham tDCS. Another recent RCT conducted by Wang et al.¹²⁴ reported a significantly higher improvement in the swallowing function after treating with an anodal tDCS, conventional swallowing rehabilitation therapy and catheter balloon dilatation in patients with CP dysfunction after brain-stem stroke.

In tDCS, a weak direct current is used to stimulate the cerebral cortex¹²⁵. tDCS has the advantage that it is easy to apply and safe with relatively few side effects. The side effects of tDCS are mild and include local erythema, tingling, or itching. In conclusion, tDCS applied to the contralateral or bilateral hemispheres is beneficial for the improvement of swallowing function when combined with swallowing therapy in patients with dysphagia.

Recommendation

tDCS is suggested for improving the swallowing function in patients with non-progressive brain lesions.

KQ 13.2. Is rTMS effective for improving the swallowing function and diet, as well as reducing risk of aspiration pneumonia?

rTMS is one of the non-invasive brain stimulation methods that is widely used to induce changes in the activity and neuroplasticity of the brain. In stroke, interhemispheric imbalance is observed with decreased excitability of the ipsilesional hemisphere and increased excitability of the contralesional hemisphere. rTMS directly delivers cortical stimulation. Low-frequency (LF) rTMS decreases cortical excitability, while high-frequency (HF) rTMS increases it. Several studies have investigated the effects of rTMS treatment on dysphagia.

A RCT conducted by Khedr et al.¹²⁶ showed that 3 Hz bilateral rTMS (10 minutes for 5 days) led to a significantly greater improvement in the swallowing function compared with sham rTMS in patients with stroke and dysphagia. Kim et al.¹²⁷ conducted a RCT to compare HF (5 Hz) ipsilateral rTMS (20 minutes for 10 days), LF (1 Hz) ipsilateral rTMS (20 minutes for 10 days), and sham rTMS in patients with brain injury and dysphagia. The results showed that FDS and PAS scores significantly improved after LF rTMS. Lim et al.¹²⁸ reported that both rTMS and NMES were effective for improving dysphagia and no significant differences were found between rTMS and NMES. Both HF and LF rTMS (3 Hz and 1 Hz, respectively) were reported to be effective for dysphagia in a RCT by Du et al.¹²⁹. A study conducted by Park et al.¹³⁰ showed that bilateral stimulation was superior to unilateral or sham stimulation in 35 patients with stroke and dysphagia. Similarly, Zhang et al.¹³¹ explained that bilateral rTMS combined with NMES produced higher cortical excitability and better swallowing function recovery compared to unilateral or sham rTMS.

Applying unilateral (LF and HF rTMS over the unaffected and affected hemispheres, respectively) or bilateral rTMS are recommended to improve the swallowing function and induce cortical neuroplasticity in patients with dysphagia. Since rTMS is a relatively new intervention in terms of applicability, additional costs are expected in addition to existing interventions,

so cost-resource allocation needs to be considered. In addition, since rTMS treatment is not currently covered by medical insurance benefits, active attention by medical staff and institutional arrangements are needed.

Recommendation

rTMS is suggested for improving swallowing function in patients with non-progressive brain lesions.

KQ 14. Does enteral tube feeding improve the clinical course, survival, or nutritional status of patients who are likely to have oropharyngeal dysphagia for a long period of time?

Under nutrition is common in patients with dysphagia and nutritional status can deteriorate as the disease persists. There are two different methods for enteral tube feeding for patients with severe dysphagia. For nasogastric feeding, a tube is inserted through the nose to supply nutrition to the stomach. During a percutaneous endoscopic gastrostomy (PEG), a tube is inserted through a hole directly into the stomach to supply nutrients. Nutrition supply using a nasogastric tube is the most commonly used non-invasive treatment, but complications such as aspiration pneumonia, reflux esophagitis, and esophageal ulcers caused by mechanical stimulation may occur. Also, a nasogastric tube must be replaced once a month. If nasogastric tube feeding is expected to be required for a long period of time, gastrostomy is recommended. However, gastrostomy is an invasive procedure that can sometimes cause complications, such as inflammation and bleeding in the procedure site. It is still controversial whether feeding through a gastrostomy, rather than maintaining a nasogastric tube, affects the clinical course and mortality of patients.

Six RCTs investigated the effect of early or preventive enteral tube feeding, including one study on patients with stroke and five studies on patients with head and neck cancer. The usefulness of early (within 72 hours after stroke) and prophylactic enteral tube feeding was investigated in patients with oropharyngeal dysphagia due to stroke and head and neck cancer.

A RCT conducted by Dennis et al.¹³² reported that early enteral tube feeding (early defined as tube feeding within 7 days of admission) was associated with an absolute reduction in risk of death and a reduction in death or poor outcome in 741 patients with stroke. Among the five RCTs conducted on patients with head and neck cancer¹³³⁻¹³⁷, three studies assessing the survival rate did not show an association between the application of early or prophylactic enteral tube feeding and the improvement in survival rate^{134,136,137}. However, Silander et al.¹³⁴ and Salas et al.¹³³ reported that prophylactic PEG was associated with significantly fewer malnourished patients over time and improved quality of life at 6 months. These studies showed that early enteral tube feeding showed overall improvement in the clinical course or survival rate in patients with oropharyngeal dysphagia.

Five RCTs studies investigated whether to supply nutrients through nasogastric and gastrostomy tubes when tube feeding is to be maintained for a long period of time¹³⁸⁻¹⁴². In most studies, patients who underwent gastrostomy showed better results in nutritional status, including blood albumin and body weight. A higher frequency of gastrointestinal bleeding, pneumonia, or urinary tract infection was reported in the nasogastric tube group compared to the gastrostomy group¹⁴⁰. Gastrostomy feeding showed a positive effect on nutritional status and weight gain compared to nasogastric feeding. Gastrostomy tube feeding is likely to bring more benefits than nasogastric tube feeding when enteral tube feeding is required for a long period of time. Therefore, switching to a gastrostomy tube is recommended in patients who need to continue nasogastric tube feeding for a long time.

Since adequate nutrition and prevention of weight loss are very important issues in clinical practice, appropriate enteral tube feeding is recommended for patients who need sufficient nutrition. If it is likely that nasogastric tube feeding will continue for a long time, selecting an appropriate candidate who would benefit from gastrostomy tube feeding is recommended.

Recommendation 14-1

In patients who are likely to have oropharyngeal dysphagia for a long period of time, early enteral tube feeding is suggested for improving the clinical course, survival rate, and neurological prognosis.

Recommendation 14-2

In patients who are likely to have oropharyngeal dysphagia for a long period of time, a gastrostomy tube feeding is suggested for improving the clinical course, survival rate, neurological prognosis, and nutritional status.

KQ 15. Does texture modification of food or liquid affect the clinical course (nutrition status or dehydration) of dysphagia?

In patients with oropharyngeal dysphagia, aspiration may occur when food is not controlled in the oral cavity during the oral phase or when an appropriate pharyngeal swallowing response is not initiated during the pharyngeal phase. If the muscle strength of the oral muscles is reduced or coordinated movements for swallowing are not appropriately performed, bolus formation or oral transit of food becomes difficult. In particular, in the case of oropharyngeal dysphagia caused by neurogenic problems, the risk of aspiration of low-viscosity food, such as water, is high due to lack of bolus control, reduced lingual propulsion, or delayed swallowing response in the pharyngeal phase¹⁴³. Aspiration occurring during swallowing can cause poor nutritional status and impede sufficient fluid intake, increasing the risk of aspiration pneumonia¹⁴⁴. Therefore, texture modification is one of the compensatory strategies for the treatment of dysphagia.

Five RCTs reported the beneficial effect of texture modification for preventing aspiration or aspiration pneumonia. A RCT conducted by Diniz et al.¹⁴⁵ reported that the use of a spoonthick consistency reduced the risk of aspiration compared with the liquid consistency in 61 patients with stroke and dysphagia. Kyodo et al.¹⁴⁶ suggested that pureed diets containing a gelling agent might reduce the risk of aspiration pneumonia possibly by decreasing pharyngeal

residues in patients with moderate to severe dysphagia.

In contrast, Robbins et al.¹⁴⁷ reported that texture modification did not affect the incidence of aspiration pneumonia. In 515 patients with dementia and Parkinson's disease who presented aspiration on a liquid diet, the 3-month cumulative incidence of pneumonia did not show any difference when texture modification (nectar or honey) was applied compared to chin-down posture. More patients who had thickened liquids showed dehydration, urinary tract infection, and fever compared to those who ingested liquid with the chin-down posture.

There were differences in disease conditions, texture modification settings, and outcome scales among the studies; however, the positive effects of texture modification on the clinical course for patients with dysphagia seem to be significant, especially for the reduction of the risk of aspiration. Texture modification is relatively easy to apply and does not require large costs; therefore, applying texture modification is recommended based on the severity of dysphagia.

Recommendation

Texture modification of food and fluids is suggested based on the severity of dysphagia to improve the clinical course (nutrition status or dehydration) of patients with oropharyngeal dysphagia.

KQ 16. Does nutrition intervention improve intake or nutritional status in patients with oropharyngeal dysphagia?

Nutrition intervention is a process related to setting specific goals to solve the patient's nutritional problems and establishing and executing plans for nutrition management. It aims to improve the quality of life by inducing changes in eating behavior and improving the patient's nutritional status¹⁴⁸. Patients with stroke often show malnutrition after hospitalization, which is presented by a reduction in muscle mass, low body mass, and low serum protein levels. Impaired oral function and dysphagia are associated with decreased oral intake, which increases the risk of malnutrition¹⁴⁹. Nutrition intervention can reduce complication rates,

admission rates, length of hospital stay, cost of care, and mortality¹⁵⁰. Thus, a multidisciplinary team approach addressing nutritional problems can help patients with oropharyngeal dysphagia.

Eight studies^{149,151-157} assessing the effectiveness of nutrition intervention in patients with dysphagia, including three RCTs and five non-randomized intervention studies, were found. A RCT conducted by Germain et al.¹⁵¹ showed that older patients who received a dysphagia-specific nutrition care program showed significant differences in weight and calorie and protein intake compared to the control group. Reyes-Torres et al.¹⁵² also reported that patients with dysphagia showed increased body weight, consumption of energy and protein, and handgrip strength after a 12-week nutrition intervention (modified consistency diet with a nectar or pudding viscosity) compared to the control group. Another RCT conducted by Taylor and Barr¹⁵³ showed that nutrition intervention (small and frequent meals) was associated with increased fluid intake. Five more observational studies showed some or no statistically significant differences in intake and nutritional status after a nutrition intervention; however, overall, nutrition intervention seems positively affect patients with dysphagia.

Recommendation

Nutrition intervention is suggested for improving intake or nutritional status in patients with oropharyngeal dysphagia.

KQ 17. Are the incidence and mortality rates of aspiration pneumonia higher in patients with oropharyngeal dysphagia compared with those without oropharyngeal dysphagia?

Pneumonia is the third leading cause of death in all age groups in Korea, and it is continuously increasing with aging¹⁵⁸. Aspiration pneumonia is a bacterial pneumonia caused by aspiration of the contents of the oropharynx or upper gastrointestinal tract colonized by pathogenic bacteria and is more severe than non-aspiration pneumonia. It has been

reported that aspiration pneumonia accounts for approximately 14.2% of community-acquired pneumonia; it is more common in older adults, requires more frequent ICU treatment, and has a longer hospital stay compared to non-aspiration pneumonia¹⁵⁹. A major risk factor for aspiration pneumonia is known as dysphagia. Dysphagia is frequently observed in vulnerable patient groups, such as older adults and patients with neurological diseases, and is associated with an increase in mortality rate, hospitalization period, and medical costs¹⁵⁹⁻¹⁶³. Therefore, it is necessary to identify the relationship between dysphagia and aspiration pneumonia and diagnose and monitor high-risk groups to provide appropriate treatment quickly and accurately.

According to the study by Lo et al.¹⁶⁴, which compared 6,979 newly diagnosed patients with dysphagia and 20,937 undiagnosed patients, the incidence of aspiration pneumonia (1.75% vs. 0.92%, $P < 0.0001$) and mortality (23.83% vs. 13.39%, $P < 0.001$) was higher in the dysphagia group than in the control group. The incidence of 1-, 3-, and 5-year aspiration pneumonia and 1-, 3-, and 5-year mortality rates after stroke diagnosis was also higher in patients with dysphagia compared to the control group¹⁶⁰. Another study also reported that among 9,930 adults aged 65 years or older who were admitted to 1,121 facilities, the presence and severity of dysphagia were related to the occurrence of aspiration pneumonia. Patients diagnosed with aspiration pneumonia had a statistically significant decline in swallowing function within 3 months compared to patients without a diagnosis (32.8% vs. 5.7%, $P < 0.001$)¹⁶⁵. In addition, other studies reported that the presence and severity of dysphagia were reported to be related to the occurrence of aspiration pneumonia¹⁶⁶⁻¹⁶⁸.

In summary, oropharyngeal dysphagia increases the risk of aspiration, and patients with oropharyngeal dysphagia have a higher incidence and mortality of aspiration pneumonia than those without dysphagia.

Recommendation

The incidence and mortality of aspiration pneumonia are higher in patients with oropharyngeal dysphagia than in those without dysphagia. Therefore, patients with dysphagia should watch out for the occurrence of pneumonia.

KQ 18. Is the oral care program effective for improving oral health and food intake?

Clean hygienic conditions in the oral cavity prevent dental caries and dryness and improve oral health and swallowing by improving oral sensation and salivation. Normal oral swallowing function can prevent swallowing problems at the pharynx stage by properly pulverizing and mixing food and saliva to form an appropriate mass that passes into the pharynx^{169,170}.

A RCT conducted by Chen et al.¹⁷¹ showed that patients who received the oral care program showed significant improvement in the oral health assessment tool but not in FOIS compared with patients who had usual oral care. Chipps et al.¹⁷² also conducted a RCT and showed that bacteria colonization decreased after the oral care program. Only two studies were found on the effectiveness of the oral care program; however, overall, it seems that there are no potential risks or side effects in the process of the oral care program, and it is easy to perform. Applying the oral care program can reduce bacterial colonization in the oral cavity and improve oral health and swallowing function.

Recommendation

The oral health care program is recommended for improving oral health and food intake in patients with oropharyngeal dysphagia.

KQ 19. Is the multidisciplinary team approach (doctor, nurse, therapist, nurse, etc.) effective for reducing complications (such as mortality, pneumonia, and other respiratory infections) in patients with oropharyngeal dysphagia?

The multidisciplinary rehabilitation team is made

up of health care professionals who work collaboratively with patients and caregivers to evaluate and manage dysphagia. In the multidisciplinary team, doctors, speech-language pathologists, nutritionists, and nurses are usually involved. Prosthodontists and dentists can sometimes also be involved in identifying structural problems related to swallowing, and social workers can be involved in discharge plans. The goal of the multidisciplinary team approach is to identify patients at risk for dysphagia, evaluate the severity of dysphagia, and provide appropriate treatment¹⁷³⁻¹⁷⁵.

A RCT conducted by Zheng et al.¹⁷⁶ showed that swallowing function significantly improved in patients with acute stroke who received a multidisciplinary team approach compared to the controls who received the conventional treatment. In four observational studies, patients who received care via the multidisciplinary team approach showed a significantly lower risk of dysphagia, pneumonia, and requirement for respiratory support, as well as greater patient satisfaction regarding swallowing function than patients who received conventional treatment^{149,177-179}.

Although the level of evidence is low, the multidisciplinary team approach should be performed because it can contribute to the improvement of swallowing function and reduce the occurrence of pneumonia. Overall, the multidisciplinary team approach seems to be effective in reducing the incidence of pneumonia during rehabilitation treatment of dysphagia.

Recommendation

A multidisciplinary team approach (doctors, nurses, therapists, etc.) is suggested for preventing complications (such as mortality, pneumonia, and other respiratory infections) in patients with oropharyngeal dysphagia.

DISCUSSION

This CPG aimed to review the literature and provide evidence-based guidelines for the assessment

and management of dysphagia. This guideline is intended to help medical staff and related subjects in charge of patients with oropharyngeal dysphagia to make safer and more effective decisions by presenting assessment and treatment guidelines based on scientific and objective evidence. Therefore, this guideline aims to effectively deliver the information necessary for decision-making and provide patients with dysphagia with appropriate education, evaluation, and treatment. This will enable patients to improve their symptoms of dysphagia and improve their quality of life. The ultimate goal is to reduce complications and mortality due to the presence of dysphagia.

First, the effectiveness of diagnostic testing for oropharyngeal dysphagia was assessed. Early screening was effective in patients with suspected oropharyngeal dysphagia to reduce the occurrence of pneumonia with a high level of evidence. The recommendation levels were strong, and we concluded that if patients with oropharyngeal dysphagia are screened early and are provided with appropriate treatment as early as possible, serious complications, such as pneumonia, can be prevented. Standardized screening tests, such as BDST, GUSS, SSA, TOR-BSST, or Clinical Functional Scale for Dysphagia, were effective in diagnosing dysphagia, as well as single screening tests, such as the 3-oz water swallow test or the volume-viscosity swallow test. For high-risk patients who are unable to swallow their saliva properly, a standardized screening test was safer than a single screening test, which requires swallowing water. A standardized screening test tool can evaluate a patient's swallowing function without directly swallowing food, and it was recommended to diagnose dysphagia in patients with suspected oropharyngeal dysphagia. To visualize a series of swallowing processes occurring in the oral, pharyngeal, and esophageal phases, and detect dysphagia, VFSS was strongly recommended for diagnosis of dysphagia with moderate levels of evidence. FEES can also be used to visualize the swallowing function, but FEES does not seem to be more effective than VFSS. Considering the benefits of

the two diagnostic methods, we concluded that the two test methods complement each other and can be performed together.

There are various treatment methods for managing dysphagia. We investigated the efficacy of various rehabilitative strategies that are commonly used for the treatment of dysphagia. Oropharyngeal sensory stimulation therapy was effective in improving swallowing function and quality of life, as well as tongue and pharyngeal muscle strengthening exercises, which improved the swallowing function and reduced aspiration pneumonia in patients with brain lesions and head and neck tumors. EMSTs were also effective in improving the swallowing function and quality of life. Compensatory maneuvers, including chin tuck, chin down, and effortful swallow, were useful for improving the swallowing function and were recommended as effective rehabilitative techniques for dysphagia management. The combination of NMES and swallowing therapy was also recommended for improving the swallowing function in patients with non-progressive neuropathic lesions and head and neck cancer. The efficacy of PES for improving swallowing function and preventing pneumonia was unclear.

Treating dysphagia with drugs by stimulating the TRP channel, such as capsaicin, menthol, or piperine, was effective for improving the swallowing function, although it is well applied in clinical practice. Thus, future studies should investigate the effect of drug treatment. In addition, swallowing treatment using biofeedback as an adjunct to conventional swallowing therapy was effective in improving swallowing function. CP botulinum toxin injection, CP myotomy, and balloon dilatation seem to be effective treatment options for patients with CP dysfunction but should be performed after considering the advantages and disadvantages of the treatments. The effect of non-invasive brain stimulation treatment methods, such as tDCS and rTMS, were also investigated, and both of them were effective in improving the swallowing function in patients with non-progressive brain lesions.

As for the nutrition issues, we concluded that early enteral tube feeding improved the clinical course, survival rate, and neurological prognosis for patients who are likely to have oropharyngeal dysphagia for a prolonged period. In addition, gastrostomy tube feeding should also be considered to improve the clinical course, survival rate, neurological prognosis, and nutritional status in patients who are likely to have oropharyngeal dysphagia for an extended period. In addition, texture modification of food and fluids based on the severity of dysphagia improved the clinical course of patients with oropharyngeal dysphagia, especially in reducing the risk of aspiration. Nutrition intervention, including addressing nutritional problems and establishing plans for nutrition management, was recommended because a literature search showed that it helps improve intake or nutritional status in patients with oropharyngeal dysphagia.

Other issues associated with dysphagia have also been addressed in this CPG. The current evidence showed that the incidence and mortality of aspiration pneumonia were, in fact, higher in patients with oropharyngeal dysphagia than those without dysphagia. Providing the oral health care program improved oral health and food intake in patients with oropharyngeal dysphagia. Additionally, a multidisciplinary team approach, including doctors in various fields, nurses, and therapists, was effective in preventing complications (such as mortality, pneumonia, and other respiratory infections). Various methods for the treatment of dysphagia should be performed, considering various aspects in patients with oropharyngeal dysphagia.

This CPG has several limitations. First, the statistical significance was not evaluated, and meta-analysis was not performed. Second, although the level of evidence for each clinical question was established based on the results of studies abroad, the recommendations of this CPG were primarily based on the applicability of the resources and healthcare system in Korea.

In conclusion, this CPG is the first guideline that

provides the levels of evidence of relevant literature and the consensus of multidisciplinary experts regarding issues related to oropharyngeal dysphagia. Physicians, patients, caregivers, and other healthcare professionals are expected to widely read this CPG to improve their understanding and treatment of dysphagia.

CONFLICTS OF INTEREST

The members who were involved in this guideline had no other conflicts of interest (COI). The COI was required to determine whether or not these persons should be involved in the development of similar guidelines, employment, financial interests, and other potential interests. No potential conflicts of interest relevant to this article were reported.

ACKNOWLEDGEMENTS

This guideline was developed with financial support from the Korean Academy of Rehabilitation Medicine and the Korean Dysphagia Society. The development of this CPG was not influenced by the supporting academies and was not supported by other groups.

SUPPLEMENTARY MATERIALS

Supplementary materials can be found via <https://doi.org/10.34160/jkds.23.010>.

REFERENCES

1. Kang TW, Kim BR. The effects of head-lift exercise on swallowing function in patients with stroke. *J Korean Soc Phys Med*. 2018;13:89-95. <https://doi.org/10.13066/kspm.2018.13.2.89>
2. Malagelada JR, Bazzoli F, Boeckxstaens G, De Looze D, Fried M, Kahrilas P, et al. World gastroenterology organisation global guidelines: dysphagia--global guidelines and cascades update September 2014. *J Clin Gastroenterol*. 2015;49:370-8. <https://doi.org/10.1097/mcg.0000000000000307>
3. Lacy BE, Paquette L, Robertson DJ, Kelley ML Jr, Weiss JE. The clinical utility of esophageal manometry. *J Clin Gastroenterol*. 2009;43:809-15. <https://doi.org/10.1097/mcg.0b013e31818ddbd5>
4. Cichero JA, Lam P, Steele CM, Hanson B, Chen J, Dantas RO, et al. Development of international terminology and definitions for texture-modified foods and thickened fluids used in dysphagia management: the IDDSI framework. *Dysphagia*. 2017;32:293-314. <https://doi.org/10.1007/s00455-016-9758-y>
5. Baijens LW, Clavé P, Cras P, Ekberg O, Forster A, Kolb GF, et al. European Society for Swallowing Disorders - European Union Geriatric Medicine Society white paper: oropharyngeal dysphagia as a geriatric syndrome. *Clin Interv Aging*. 2016;11:1403-28. <https://doi.org/10.2147/cia.s107750>
6. Smith R, Bryant L, Hemsley B. The true cost of dysphagia on quality of life: the views of adults with swallowing disability. *Int J Lang Commun Disord*. 2023;58:451-66. <https://doi.org/10.1111/1460-6984.12804>
7. Wu XS, Miles A, Braakhuis A. Nutritional intake and meal composition of patients consuming texture modified diets and thickened fluids: a systematic review and meta-analysis. *Healthcare (Basel)*. 2020;8:579. <https://doi.org/10.3390/healthcare8040579>
8. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. <https://doi.org/10.1136/bmj.d5928>
9. Kim SY, Park JE, Lee YJ, Seo HJ, Sheen SS, Hahn S, et al. Testing a tool for assessing the risk of bias for non-randomized studies showed moderate reliability and promising validity. *J Clin Epidemiol*. 2013;66:408-14. <https://doi.org/10.1016/j.jclinepi.2012.09.016>
10. Shea BJ, Hamel C, Wells GA, Bouter LM, Kristjansson E, Grimshaw J, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol*. 2009;62:1013-20. <https://doi.org/10.1016/j.jclinepi.2008.10.009>
11. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924-6. <https://doi.org/10.1136/bmj.39489.470347.ad>
12. Schmidt Leuenberger JM, Hokscho B, Luder G, Schmid RA, Verra ML, Dorn P. Early assessment and management of dysphagia after lung resection: a randomized controlled trial. *Ann Thorac Surg*. 2019;108:1059-64.
13. Posillico SE, Golob JF, Rinker AD, Kreiner LA, West RS, Conrad-Schnetz KJ, et al. Bedside dysphagia screens in patients with traumatic cervical injuries: an ideal tool for an under-recognized problem. *J Trauma Acute Care Surg*. 2018;85:697-703. <https://doi.org/10.1097/ta.0000000000002035>
14. Wangen T, Hatlevig J, Pifer G, Vitale K. Preventing aspiration complications: implementing a swallow screening tool. *Clin Nurse Spec*. 2019;33:237-43.

- <https://doi.org/10.1097/nur.0000000000000471>
15. Hinchey JA, Shephard T, Furie K, Smith D, Wang D, Tonn S. Formal dysphagia screening protocols prevent pneumonia. *Stroke*. 2005;36:1972-6.
<https://doi.org/10.1161/01.str.0000177529.86868.8d>
 16. Yeh SJ, Huang KY, Wang TG, Chen YC, Chen CH, Tang SC, et al. Dysphagia screening decreases pneumonia in acute stroke patients admitted to the stroke intensive care unit. *J Neurol Sci*. 2011;306:38-41.
<https://doi.org/10.1016/j.jns.2011.04.001>
 17. Sørensen RT, Rasmussen RS, Overgaard K, Lerche A, Johansen AM, Lindhardt T. Dysphagia screening and intensified oral hygiene reduce pneumonia after stroke. *J Neurosci Nurs*. 2013;45:139-46.
<https://doi.org/10.1097/jnn.0b013e31828a412c>
 18. Titworth WL, Abram J, Fullerton A, Hester J, Guin P, Waters MF, et al. Prospective quality initiative to maximize dysphagia screening reduces hospital-acquired pneumonia prevalence in patients with stroke. *Stroke*. 2013;44:3154-60.
<https://doi.org/10.1161/strokeaha.111.000204>
 19. Schrock JW, Lou L, Ball BAW, Van Etten J. The use of an emergency department dysphagia screen is associated with decreased pneumonia in acute strokes. *Am J Emerg Med*. 2018;36:2152-4.
<https://doi.org/10.1016/j.ajem.2018.03.046>
 20. Teuschl Y, Trapl M, Ratajczak P, Matz K, Dachenhausen A, Brainin M. Systematic dysphagia screening and dietary modifications to reduce stroke-associated pneumonia rates in a stroke-unit. *PLoS One*. 2018;13:e0192142.
<https://doi.org/10.1371/journal.pone.0192142>
 21. Taveira I, Silva S, Bonança Í, Parreira D, Antunes C. Recognizing dysphagia: implementation of an in-hospital screening protocol. *Ir J Med Sci*. 2021;190:605-8.
<https://doi.org/10.1007/s11845-020-02371-8>
 22. Hammond CAS, Goldstein LB. Cough and aspiration of food and liquids due to oral-pharyngeal dysphagia: ACCP evidence-based clinical practice guidelines. *Chest*. 2006;129(1 Suppl):154S-68S.
https://doi.org/10.1378/chest.129.1_suppl.154s
 23. Shin JC, Kim DY, Heo JH, Song W, Yi SH. Comparison between Gugging Swallowing Screen and other dysphagia screening tests. *Brain Neurorehabil*. 2009;2:146-54.
 24. DePippo KL, Holas MA, Reding MJ. The Burke dysphagia screening test: validation of its use in patients with stroke. *Arch Phys Med Rehabil*. 1994;75:1284-6.
 25. Perry L. Screening swallowing function of patients with acute stroke. Part one: Identification, implementation and initial evaluation of a screening tool for use by nurses. *J Clin Nurs*. 2001;10:463-73.
<https://doi.org/10.1046/j.1365-2702.2001.00501.x>
 26. Trapl M, Enderle P, Nowotny M, Teuschl Y, Matz K, Dachenhausen A, et al. Dysphagia bedside screening for acute-stroke patients: the Gugging Swallowing Screen. *Stroke*. 2007;38:2948-52.
<https://doi.org/10.1161/strokeaha.107.483933>
 27. Lopes M, Freitas E, Oliveira M, Dantas E, Azevedo N, Rodrigues P, et al. Impact of the systematic use of the Gugging Swallowing Screen in patients with acute ischaemic stroke. *Eur J Neurol*. 2019;26:722-6.
<https://doi.org/10.1111/ene.13825>
 28. Rugiu MG. Role of videofluoroscopy in evaluation of neurologic dysphagia. *Acta Otorhinolaryngol Ital*. 2007;27:306-16.
 29. Doggett DL, Tappe KA, Mitchell MD, Chapell R, Coates V, Turkelson CM. Prevention of pneumonia in elderly stroke patients by systematic diagnosis and treatment of dysphagia: an evidence-based comprehensive analysis of the literature. *Dysphagia*. 2001;16:279-95.
<https://doi.org/10.1007/s00455-001-0087-3>
 30. Perry L, Love CP. Screening for dysphagia and aspiration in acute stroke: a systematic review. *Dysphagia*. 2001;16:7-18.
<https://doi.org/10.1007/pl00021290>
 31. Shem KL, Castillo K, Wong SL, Chang J, Kao MC, Kolakowsky-Hayner SA. Diagnostic accuracy of bedside swallow evaluation versus videofluoroscopy to assess dysphagia in individuals with tetraplegia. *PM R*. 2012;4:283-9.
<https://doi.org/10.1016/j.pmrj.2012.01.002>
 32. Wilson RD, Howe EC. A cost-effectiveness analysis of screening methods for dysphagia after stroke. *PM R*. 2012;4:273-82.
<https://doi.org/10.1016/j.pmrj.2011.09.006>
 33. Leigh J, Lim JY, Han MK, Bae HJ, Kim WS, Paik NJ. A prospective comparison between bedside swallowing screening test and videofluoroscopic swallowing study in post-stroke dysphagia. *Brain Neurorehabil*. 2016;9:e7.
<https://doi.org/10.12786/bn.2016.9.e7>
 34. Rangarathnam B, McCullough GH. Utility of a clinical swallowing exam for understanding swallowing physiology. *Dysphagia*. 2016;31:491-7.
<https://doi.org/10.1007/s00455-016-9702-1>
 35. Aviv JE. Prospective, randomized outcome study of endoscopy versus modified barium swallow in patients with dysphagia. *Laryngoscope*. 2000;110:563-74.
<https://doi.org/10.1097/00005537-200004000-00008>
 36. Wu CH, Hsiao TY, Chen JC, Chang YC, Lee SY. Evaluation of swallowing safety with fiberoptic endoscope: comparison with videofluoroscopic technique. *Laryngoscope*. 1997;107:396-401.
<https://doi.org/10.1097/00005537-199703000-00023>
 37. Fattori B, Giusti P, Mancini V, Grosso M, Barillari MR, Bastiani L, et al. Comparison between videofluoroscopy, fiberoptic endoscopy and scintigraphy for diagnosis of oro-pharyngeal dysphagia. *Acta Otorhinolaryngol Ital*. 2016;36:395-402.
<https://doi.org/10.14639/0392-100x-829>
 38. Kelly AM, Leslie P, Beale T, Payten C, Drinnan MJ. Fiberoptic endoscopic evaluation of swallowing and videofluoroscopy: does examination type influence perception of pharyngeal residue severity? *Clin Otolaryngol*. 2006;31:425-32.

- <https://doi.org/10.1111/j.1749-4486.2006.01292.x>
39. Kelly AM, Drinnan MJ, Leslie P. Assessing penetration and aspiration: how do videofluoroscopy and fiberoptic endoscopic evaluation of swallowing compare? *Laryngoscope*. 2007;117:1723-7.
<https://doi.org/10.1097/mlg.0b013e318123ee6a>
 40. Power M, Fraser C, Hobson A, Rothwell JC, Mistry S, Nicholson DA, et al. Changes in pharyngeal corticobulbar excitability and swallowing behavior after oral stimulation. *Am J Physiol Gastrointest Liver Physiol*. 2004;286:G45-50.
<https://doi.org/10.1152/ajpgi.00114.2003>
 41. Maeda K, Koga T, Akagi J. Interferential current sensory stimulation, through the neck skin, improves airway defense and oral nutrition intake in patients with dysphagia: a double-blind randomized controlled trial. *Clin Interv Aging*. 2017;12:1879-86.
<https://doi.org/10.2147/cia.s140746>
 42. Zhang M, Tao T, Zhang ZB, Zhu X, Fan WG, Pu LJ, et al. Effectiveness of neuromuscular electrical stimulation on patients with dysphagia with medullary infarction. *Arch Phys Med Rehabil*. 2016;97:355-62.
<https://doi.org/10.1016/j.apmr.2015.10.104>
 43. Rofes L, Arreola V, López I, Martín A, Sebastián M, Ciurana A, et al. Effect of surface sensory and motor electrical stimulation on chronic poststroke oropharyngeal dysfunction. *Neurogastroenterol Motil*. 2013;25:888-e701.
<https://doi.org/10.1111/nmo.12211>
 44. Lazarus CL, Husaini H, Falciglia D, DeLacure M, Branski RC, Kraus D, et al. Effects of exercise on swallowing and tongue strength in patients with oral and oropharyngeal cancer treated with primary radiotherapy with or without chemotherapy. *Int J Oral Maxillofac Surg*. 2014;43:523-30.
<https://doi.org/10.1016/j.ijom.2013.10.023>
 45. Mepani R, Antonik S, Massey B, Kern M, Logemann J, Pauloski B, et al. Augmentation of deglutitive thyrohyoid muscle shortening by the Shaker Exercise. *Dysphagia*. 2009;24:26-31.
<https://doi.org/10.1007/s00455-008-9167-y>
 46. McCullough GH, Kamarunas E, Mann GC, Schmidley JW, Robbins JA, Crary MA. Effects of Mendelsohn maneuver on measures of swallowing duration post stroke. *Top Stroke Rehabil*. 2012;19:234-43.
<https://doi.org/10.1310/tsr1903-234>
 47. McCullough GH, Kim Y. Effects of the Mendelsohn maneuver on extent of hyoid movement and UES opening post-stroke. *Dysphagia*. 2013;28:511-9.
<https://doi.org/10.1007/s00455-013-9461-1>
 48. Kotz T, Federman AD, Kao J, Milman L, Packer S, Lopez-Prieto C, et al. Prophylactic swallowing exercises in patients with head and neck cancer undergoing chemoradiation: a randomized trial. *Arch Otolaryngol Head Neck Surg*. 2012;138:376-82.
<https://doi.org/10.1001/archoto.2012.187>
 49. Eom MJ, Chang MY, Oh DH, Kim HD, Han NM, Park JS. Effects of resistance expiratory muscle strength training in elderly patients with dysphagic stroke. *NeuroRehabilitation*. 2017;41:747-52.
<https://doi.org/10.3233/nre-172192>
 50. Moon JH, Jung JH, Won YS, Cho HY, Cho K. Effects of expiratory muscle strength training on swallowing function in acute stroke patients with dysphagia. *J Phys Ther Sci*. 2017;29:609-12.
<https://doi.org/10.1589/jpts.29.609>
 51. Park JS, Oh DH, Chang MY, Kim KM. Effects of expiratory muscle strength training on oropharyngeal dysphagia in subacute stroke patients: a randomised controlled trial. *J Oral Rehabil*. 2016;43:364-72.
<https://doi.org/10.1111/joor.12382>
 52. Silverman EP, Miller S, Zhang Y, Hoffman-Ruddy B, Yeager J, Daly JJ. Effects of expiratory muscle strength training on maximal respiratory pressure and swallow-related quality of life in individuals with multiple sclerosis. *Mult Scler J Exp Transl Clin*. 2017;3:2055217317710829.
<https://doi.org/10.1177/2055217317710829>
 53. Solazzo A, Monaco L, Del Vecchio L, Tamburrini S, Iacobellis F, Berritto D, et al. Investigation of compensatory postures with videofluoromanometry in dysphagia patients. *World J Gastroenterol*. 2012;18:2973-8.
<https://doi.org/10.3748/wjg.v18.i23.2973>
 54. Ra JY, Hyun JK, Ko KR, Lee SJ. Chin tuck for prevention of aspiration: effectiveness and appropriate posture. *Dysphagia*. 2014;29:603-9.
<https://doi.org/10.1007/s00455-014-9551-8>
 55. Miyamoto T, Kumai Y, Matsubara K, Kodama N, Satoh C, Orita Y. Different types of dysphagia alleviated by the chin-down position. *Auris Nasus Larynx*. 2021;48:928-33.
<https://doi.org/10.1016/j.anl.2021.02.008>
 56. Logemann JA, Pauloski BR, Rademaker AW, Colangelo LA. Super-supraglottic swallow in irradiated head and neck cancer patients. *Head Neck*. 1997;19:535-40.
[https://doi.org/10.1002/\(sici\)1097-0347\(199709\)19:6<3C535::aid-hed11%3E3.0.co;2-4](https://doi.org/10.1002/(sici)1097-0347(199709)19:6<3C535::aid-hed11%3E3.0.co;2-4)
 57. Bülow M, Olsson R, Ekberg O. Supraglottic swallow, effortful swallow, and chin tuck did not alter hypopharyngeal intrabolus pressure in patients with pharyngeal dysfunction. *Dysphagia*. 2002;17:197-201.
<https://doi.org/10.1007/s00455-002-0050-y>
 58. González-Fernández M, Ottenstein L, Atanelov L, Christian AB. Dysphagia after stroke: an overview. *Curr Phys Med Rehabil Rep*. 2013;1:187-96.
<https://doi.org/10.1007/s40141-013-0017-y>
 59. Carnaby GD, LaGorio L, Silliman S, Crary M. Exercise-based swallowing intervention (McNeill Dysphagia Therapy) with adjunctive NMES to treat dysphagia post-stroke: a double-blind placebo-controlled trial. *J Oral Rehabil*. 2020;47:501-10.
<https://doi.org/10.1111/joor.12928>

60. Guillén-Solà A, Messagi Sartor M, Bofill Soler N, Duarte E, Barrera MC, Marco E. Respiratory muscle strength training and neuromuscular electrical stimulation in subacute dysphagic stroke patients: a randomized controlled trial. *Clin Rehabil*. 2017;31:761-71. <https://doi.org/10.1177/0269215516652446>
61. Konecny P, Elfmark M. Electrical stimulation of hyoid muscles in post-stroke dysphagia. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2018;162:40-2. <https://doi.org/10.5507/bp.2017.043>
62. Lee KW, Kim SB, Lee JH, Lee SJ, Ri JW, Park JG. The effect of early neuromuscular electrical stimulation therapy in acute/subacute ischemic stroke patients with Dysphagia. *Ann Rehabil Med*. 2014;38:153-9. <https://doi.org/10.5535/arm.2014.38.2.153>
63. Lim KB, Lee HJ, Lim SS, Choi YI. Neuromuscular electrical and thermal-tactile stimulation for dysphagia caused by stroke: a randomized controlled trial. *J Rehabil Med*. 2009;41:174-8. <https://doi.org/10.2340/16501977-0317>
64. Jung SH, Kim YA, Hwang NK, Park JS, Kim YH. Effects of neuromuscular electrical stimulation in combination with saliva or dry swallowing in stroke patients with dysphagia. *J Korean Dysphagia Soc*. 2018;8:103-9.
65. Sproson L, Pownall S, Enderby P, Freeman J. Combined electrical stimulation and exercise for swallow rehabilitation post-stroke: a pilot randomized control trial. *Int J Lang Commun Disord*. 2018;53:405-17. <https://doi.org/10.1111/1460-6984.12359>
66. Terré R, Mearin F. A randomized controlled study of neuromuscular electrical stimulation in oropharyngeal dysphagia secondary to acquired brain injury. *Eur J Neurol*. 2015;22:687-e44. <https://doi.org/10.1111/ene.12631>
67. Xia W, Zheng C, Lei Q, Tang Z, Hua Q, Zhang Y, et al. Treatment of post-stroke dysphagia by Vitalstim therapy coupled with conventional swallowing training. *J Huazhong Univ Sci Technolog Med Sci*. 2011;31:73-6. <https://doi.org/10.1007/s11596-011-0153-5>
68. Kushner DS, Peters K, Eroglu ST, Perless-Carroll M, Johnson-Greene D. Neuromuscular electrical stimulation efficacy in acute stroke feeding tube-dependent dysphagia during inpatient rehabilitation. *Am J Phys Med Rehabil*. 2013;92:486-95. <https://doi.org/10.1097/phm.0b013e31828762ec>
69. Toyama K, Matsumoto S, Kurasawa M, Setoguchi H, Noma T, Takenaka K, et al. Novel neuromuscular electrical stimulation system for treatment of dysphagia after brain injury. *Neurol Med Chir (Tokyo)*. 2014;54:521-8. <https://doi.org/10.2176/nmc.oa.2013-0341>
70. Pauloski BR. Rehabilitation of dysphagia following head and neck cancer. *Phys Med Rehabil Clin N Am*. 2008;19:889-928, x. <https://doi.org/10.1016/j.pmr.2008.05.010>
71. Ryu JS, Kang JY, Park JY, Nam SY, Choi SH, Roh JL, et al. The effect of electrical stimulation therapy on dysphagia following treatment for head and neck cancer. *Oral Oncol*. 2009;45:665-8. <https://doi.org/10.1016/j.oraloncology.2008.10.005>
72. Bhatt AD, Goodwin N, Cash E, Bhatt G, Silverman CL, Spanos WJ, et al. Impact of transcutaneous neuromuscular electrical stimulation on dysphagia in patients with head and neck cancer treated with definitive chemoradiation. *Head Neck*. 2015;37:1051-6. <https://doi.org/10.1002/hed.23708>
73. Lin PH, Hsiao TY, Chang YC, Ting LL, Chen WS, Chen SC, et al. Effects of functional electrical stimulation on dysphagia caused by radiation therapy in patients with nasopharyngeal carcinoma. *Support Care Cancer*. 2011;19:91-9. <https://doi.org/10.1007/s00520-009-0792-2>
74. Dziewas R, Stellato R, van der Tweel I, Walther E, Werner CJ, Braun T, et al. Pharyngeal electrical stimulation for early decannulation in tracheotomised patients with neurogenic dysphagia after stroke (PHAST-TRAC): a prospective, single-blinded, randomised trial. *Lancet Neurol*. 2018;17:849-59. [https://doi.org/10.1016/s1474-4422\(18\)30255-2](https://doi.org/10.1016/s1474-4422(18)30255-2)
75. Bath PM, Scutt P, Love J, Clavé P, Cohen D, Dziewas R, et al. Pharyngeal electrical stimulation for treatment of dysphagia in subacute stroke: a randomized controlled trial. *Stroke*. 2016;47:1562-70. <https://doi.org/10.1161/strokeaha.115.012455>
76. Jayasekaran V, Singh S, Tyrrell P, Michou E, Jefferson S, Mistry S, et al. Adjunctive functional pharyngeal electrical stimulation reverses swallowing disability after brain lesions. *Gastroenterology*. 2010;138:1737-46. <https://doi.org/10.1053/j.gastro.2010.01.052>
77. Suntrup S, Marian T, Schröder JB, Suttrup I, Muhle P, Oelenberg S, et al. Electrical pharyngeal stimulation for dysphagia treatment in tracheotomized stroke patients: a randomized controlled trial. *Intensive Care Med*. 2015;41:1629-37. <https://doi.org/10.1007/s00134-015-3897-8>
78. Vasant DH, Michou E, O'Leary N, Vail A, Mistry S, Hamdy S. Pharyngeal electrical stimulation in dysphagia poststroke: a prospective, randomized single-blinded interventional study. *Neurorehabil Neural Repair*. 2016;30:866-75. <https://doi.org/10.1177/1545968316639129>
79. Ebihara T, Takahashi H, Ebihara S, Okazaki T, Sasaki T, Watando A, et al. Capsaicin troche for swallowing dysfunction in older people. *J Am Geriatr Soc*. 2005;53:824-8. <https://doi.org/10.1111/j.1532-5415.2005.53261.x>
80. Ortega O, Rofes L, Martin A, Arreola V, López I, Clavé P. A comparative study between two sensory stimulation strategies after two weeks treatment on older patients with oropharyngeal dysphagia. *Dysphagia*. 2016;31:706-16. <https://doi.org/10.1007/s00455-016-9736-4>
81. Rofes L, Arreola V, Martin A, Clavé P. Natural cap-

- saicinoids improve swallow response in older patients with oropharyngeal dysphagia. *Gut*. 2013;62:1280-7. <https://doi.org/10.1136/gutjnl-2011-300753>
82. Cui F, Yin Q, Wu C, Shen M, Zhang Y, Ma C, et al. Capsaicin combined with ice stimulation improves swallowing function in patients with dysphagia after stroke: a randomised controlled trial. *J Oral Rehabil*. 2020;47:1297-303. <https://doi.org/10.1111/joor.13068>
 83. Wang Z, Wu L, Fang Q, Shen M, Zhang L, Liu X. Effects of capsaicin on swallowing function in stroke patients with dysphagia: a randomized controlled trial. *J Stroke Cerebrovasc Dis*. 2019;28:1744-51. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.02.008>
 84. Ebihara T, Ebihara S, Watando A, Okazaki T, Asada M, Ohnui T, et al. Effects of menthol on the triggering of the swallowing reflex in elderly patients with dysphagia. *Br J Clin Pharmacol*. 2006;62:369-71. <https://doi.org/10.1111/j.1365-2125.2006.02666.x>
 85. Rofes L, Arreola V, Martin A, Clavé P. Effect of oral piperine on the swallow response of patients with oropharyngeal dysphagia. *J Gastroenterol*. 2014;49:1517-23. <https://doi.org/10.1007/s00535-013-0920-0>
 86. Giggins OM, Persson UM, Caulfield B. Biofeedback in rehabilitation. *J Neuroeng Rehabil*. 2013;10:60. <https://doi.org/10.1186/1743-0003-10-60>
 87. Basmajian JV. Biofeedback: principles and practice for clinicians. Oxford: Williams & Wilkins, 1979.
 88. Shin YA, Kwon HC, Lee SM. Effects of biofeedback training on the suprahyoid muscle activity of stroke with swallowing disorder. *J Korean Soc Integr Med*. 2019;7:27-36.
 89. Moon JH, Kim GY, Won YS. Effects of swallowing training with biofeedback on swallowing function and satisfaction in acute stroke patients with dysphagia. *J Korea Contents Assoc*. 2017;17:63-71. <https://doi.org/10.5392/JKCA.2017.17.04.063>
 90. Bogaardt HC, Grolman W, Fokkens WJ. The use of biofeedback in the treatment of chronic dysphagia in stroke patients. *Folia Phoniatr Logop*. 2009;61:200-5. <https://doi.org/10.1159/000227997>
 91. Crary MA, Carnaby Mann GD, Groher ME, Helseth E. Functional benefits of dysphagia therapy using adjunctive sEMG biofeedback. *Dysphagia*. 2004;19:160-4. <https://doi.org/10.1007/s00455-004-0003-8>
 92. Li CM, Lee HY, Hsieh SH, Wang TG, Wang HP, Chen JJJ. Development of innovative feedback device for swallowing therapy. *J Med Biol Eng*. 2016;36:357-68. <https://doi.org/10.1007/s40846-016-0146-8>
 93. Li CM, Wang TG, Lee HY, Wang HP, Hsieh SH, Chou M, et al. Swallowing training combined with game-based biofeedback in poststroke dysphagia. *PM R*. 2016;8:773-9. <https://doi.org/10.1016/j.pmrj.2016.01.003>
 94. Park JS, Oh DH, Hwang NK, Lee JH. Effects of neuromuscular electrical stimulation combined with effortful swallowing on post-stroke oropharyngeal dysphagia: a randomised controlled trial. *J Oral Rehabil*. 2016;43:426-34. <https://doi.org/10.1111/joor.12390>
 95. Dewan K, Santa Maria C, Noel J. Cricopharyngeal achalasia: management and associated outcomes-a scoping review. *Otolaryngol Head Neck Surg*. 2020;163:1109-13. <https://doi.org/10.1177/0194599820931470>
 96. Kocdor P, Siegel ER, Tulunay-Ugur OE. Cricopharyngeal dysfunction: a systematic review comparing outcomes of dilatation, botulinum toxin injection, and myotomy. *Laryngoscope*. 2016;126:135-41. <https://doi.org/10.1002/lary.25447>
 97. Ashman A, Dale OT, Baldwin DL. Management of isolated cricopharyngeal dysfunction: systematic review. *J Laryngol Otol*. 2016;130:611-5. <https://doi.org/10.1017/s0022215116007994>
 98. Alfonsi E, Restivo DA, Cosentino G, De Icco R, Bertino G, Schindler A, et al. Botulinum toxin is effective in the management of neurogenic dysphagia. Clinical-electrophysiological findings and tips on safety in different neurological disorders. *Front Pharmacol*. 2017;8:80. <https://doi.org/10.3389/fphar.2017.00080>
 99. Kelly EA, Koszewski IJ, Jaradeh SS, Merati AL, Blumin JH, Bock JM. Botulinum toxin injection for the treatment of upper esophageal sphincter dysfunction. *Ann Otol Rhinol Laryngol*. 2013;122:100-8. <https://doi.org/10.1177/000348941312200205>
 100. Jeong SH, Kim YJ, Kim YJ, Park KD, Kim EJ, Chung JW, et al. Endoscopic botulinum toxin injection for treatment of pharyngeal dysphagia in patients with cricopharyngeal dysfunction. *Scand J Gastroenterol*. 2018;53:1201-5. <https://doi.org/10.1080/00365521.2018.1506820>
 101. Kim MS, Kim GW, Rho YS, Kwon KH, Chung EJ. Office-based electromyography-guided botulinum toxin injection to the cricopharyngeus muscle: optimal patient selection and technique. *Ann Otol Rhinol Laryngol*. 2017;126:349-56. <https://doi.org/10.1177/0003489416689469>
 102. Pitman M, Weissbrod P. Endoscopic CO2 laser cricopharyngeal myotomy. *Laryngoscope*. 2009;119:45-53. <https://doi.org/10.1002/lary.20032>
 103. McKenna JA, Dedo HH. Cricopharyngeal myotomy: indications and technique. *Ann Otol Rhinol Laryngol*. 1992;101:216-21. <https://doi.org/10.1177/000348949210100304>
 104. Brigand C, Ferraro P, Martin J, Duranceau A. Risk factors in patients undergoing cricopharyngeal myotomy. *Br J Surg*. 2007;94:978-83. <https://doi.org/10.1002/bjs.5760>
 105. Dauer E, Salassa J, Iuga L, Kasperbauer J. Endoscopic laser vs open approach for cricopharyngeal myotomy. *Otolaryngol Head Neck Surg*. 2006;134:830-5.

- <https://doi.org/10.1016/j.otohns.2005.12.030>
106. Takes RP, van den Hoogen FJ, Marres HA. Endoscopic myotomy of the cricopharyngeal muscle with CO₂ laser surgery. *Head Neck*. 2005;27:703-9.
<https://doi.org/10.1002/hed.20201>
 107. Ho AS, Morzaria S, Damrose EJ. Carbon dioxide laser-assisted endoscopic cricopharyngeal myotomy with primary mucosal closure. *Ann Otol Rhinol Laryngol*. 2011;120:33-9.
<https://doi.org/10.1177/000348941112000105>
 108. Lan Y, Xu G, Dou Z, Wan G, Yu F, Lin T. Biomechanical changes in the pharynx and upper esophageal sphincter after modified balloon dilatation in brainstem stroke patients with dysphagia. *Neurogastroenterol Motil*. 2013;25:e821-9.
<https://doi.org/10.1111/nmo.12209>
 109. Wei X, Yu F, Dai M, Xie C, Wan G, Wang Y, et al. Change in excitability of cortical projection after modified catheter balloon dilatation therapy in brainstem stroke patients with dysphagia: a prospective controlled study. *Dysphagia*. 2017;32:645-56.
<https://doi.org/10.1007/s00455-017-9810-6>
 110. Arenaz Búa B, Olsson R, Westin U, Rydell R, Ekberg O. Treatment of cricopharyngeal dysfunction: a comparative pilot study. *BMC Res Notes*. 2015;8:301.
<https://doi.org/10.1186/s13104-015-1266-x>
 111. Kang JH, Park RY, Lee SJ, Kim JY, Yoon SR, Jung KI. The effect of bedside exercise program on stroke patients with Dysphagia. *Ann Rehabil Med*. 2012;36:512-20.
<https://doi.org/10.5535/arm.2012.36.4.512>
 112. Cho YS, Oh DH, Paik YR, Lee JH, Park JS. Effects of bedside self-exercise on oropharyngeal swallowing function in stroke patients with dysphagia: a pilot study. *J Phys Ther Sci*. 2017;29:1815-6.
<https://doi.org/10.1589/jpts.29.1815>
 113. Mashhour K, Abdelkader R, Abdelkader L, El Hadary S, Hashem W. Swallowing exercises: will they really help head and neck cancer patients? *Asian Pac J Cancer Prev*. 2018;19:797-801.
<https://doi.org/10.22034/apjcp.2018.19.3.797>
 114. Chen SC, Huang BS, Chung CY, Lin CY, Fan KH, Chang JT, et al. Effects of a swallowing exercise education program on dysphagia-specific health-related quality of life in oral cavity cancer patients post-treatment: a randomized controlled trial. *Support Care Cancer*. 2018;26:2919-28.
<https://doi.org/10.1007/s00520-018-4148-7>
 115. Baker JM, Rorden C, Fridriksson J. Using transcranial direct-current stimulation to treat stroke patients with aphasia. *Stroke*. 2010;41:1229-36.
<https://doi.org/10.1161/strokeaha.109.576785>
 116. Jefferson S, Mistry S, Singh S, Rothwell J, Hamdy S. Characterizing the application of transcranial direct current stimulation in human pharyngeal motor cortex. *Am J Physiol Gastrointest Liver Physiol*. 2009;297:G1035-40.
<https://doi.org/10.1152/ajpgi.00294.2009>
 117. Kumar S, Wagner CW, Frayne C, Zhu L, Selim M, Feng W, et al. Noninvasive brain stimulation may improve stroke-related dysphagia: a pilot study. *Stroke*. 2011;42:1035-40.
<https://doi.org/10.1161/strokeaha.110.602128>
 118. Ahn YH, Sohn HJ, Park JS, Ahn TG, Shin YB, Park M, et al. Effect of bihemispheric anodal transcranial direct current stimulation for dysphagia in chronic stroke patients: a randomized clinical trial. *J Rehabil Med*. 2017;49:30-5.
<https://doi.org/10.2340/16501977-2170>
 119. Pingue V, Priori A, Malovini A, Pistarini C. Dual transcranial direct current stimulation for poststroke dysphagia: a randomized controlled trial. *Neurorehabil Neural Repair*. 2018;32:635-44.
<https://doi.org/10.1177/1545968318782743>
 120. Shigematsu T, Fujishima I, Ohno K. Transcranial direct current stimulation improves swallowing function in stroke patients. *Neurorehabil Neural Repair*. 2013;27:363-9.
<https://doi.org/10.1177/1545968312474116>
 121. Suntrup-Krueger S, Ringmaier C, Muhle P, Wollbrink A, Kemmling A, Hanning U, et al. Randomized trial of transcranial direct current stimulation for poststroke dysphagia. *Ann Neurol*. 2018;83:328-40.
<https://doi.org/10.1002/ana.25151>
 122. Yang EJ, Baek SR, Shin J, Lim JY, Jang HJ, Kim YK, et al. Effects of transcranial direct current stimulation (tDCS) on post-stroke dysphagia. *Restor Neurol Neurosci*. 2012;30:303-11.
<https://doi.org/10.3233/rnn-2012-110213>
 123. Sawan SAE, Reda AM, Kamel AH, Ali MAM. Transcranial direct current stimulation (tDCS): its effect on improving dysphagia in stroke patients. *Egypt J Neurol Psychiatry Neurosurg*. 2020;56:111.
<https://doi.org/10.1186/s41983-020-00246-4>
 124. Wang ZY, Chen JM, Lin ZK, Ni GX. Transcranial direct current stimulation improves the swallowing function in patients with cricopharyngeal muscle dysfunction following a brainstem stroke. *Neurol Sci*. 2020;41:569-74.
<https://doi.org/10.1007/s10072-019-04120-x>
 125. Pascual-Leone A, Valls-Solé J, Wassermann EM, Hallett M. Responses to rapid-rate transcranial magnetic stimulation of the human motor cortex. *Brain*. 1994;117(Pt 4):847-58.
<https://doi.org/10.1093/brain/117.4.847>
 126. Khedr EM, Abo-Elfetoh N, Rothwell JC. Treatment of post-stroke dysphagia with repetitive transcranial magnetic stimulation. *Acta Neurol Scand*. 2009;119:155-61.
<https://doi.org/10.1111/j.1600-0404.2008.01093.x>
 127. Kim L, Chun MH, Kim BR, Lee SJ. Effect of repetitive transcranial magnetic stimulation on patients with brain injury and Dysphagia. *Ann Rehabil Med*. 2011;35:765-71.

- <https://doi.org/10.5535/arm.2011.35.6.765>
128. Lim KB, Lee HJ, Yoo J, Kwon YG. Effect of low-frequency rTMS and NMES on subacute unilateral hemispheric stroke with dysphagia. *Ann Rehabil Med*. 2014;38:592-602.
<https://doi.org/10.5535/arm.2014.38.5.592>
 129. Du J, Yang F, Liu L, Hu J, Cai B, Liu W, et al. Repetitive transcranial magnetic stimulation for rehabilitation of poststroke dysphagia: a randomized, double-blind clinical trial. *Clin Neurophysiol*. 2016;127:1907-13.
<https://doi.org/10.1016/j.clinph.2015.11.045>
 130. Park E, Kim MS, Chang WH, Oh SM, Kim YK, Lee A, et al. Effects of bilateral repetitive transcranial magnetic stimulation on post-stroke dysphagia. *Brain Stimul*. 2017;10:75-82.
<https://doi.org/10.1016/j.brs.2016.08.005>
 131. Zhang C, Zheng X, Lu R, Yun W, Yun H, Zhou X. Repetitive transcranial magnetic stimulation in combination with neuromuscular electrical stimulation for treatment of post-stroke dysphagia. *J Int Med Res*. 2019;47:662-72.
<https://doi.org/10.1177/0300060518807340>
 132. Dennis MS, Lewis SC, Warlow C. Effect of timing and method of enteral tube feeding for dysphagic stroke patients (FOOD): a multicentre randomised controlled trial. *Lancet*. 2005;365:764-72.
[https://doi.org/10.1016/s0140-6736\(05\)17983-5](https://doi.org/10.1016/s0140-6736(05)17983-5)
 133. Salas S, Baumstarck-Barrau K, Alfonsi M, Digue L, Bagarry D, Feham N, et al. Impact of the prophylactic gastrostomy for unresectable squamous cell head and neck carcinomas treated with radio-chemotherapy on quality of life: prospective randomized trial. *Radiother Oncol*. 2009;93:503-9.
<https://doi.org/10.1016/j.radonc.2009.05.016>
 134. Silander E, Nyman J, Bove M, Johansson L, Larsson S, Hammerlid E. Impact of prophylactic percutaneous endoscopic gastrostomy on malnutrition and quality of life in patients with head and neck cancer: a randomized study. *Head Neck*. 2012;34:1-9.
<https://doi.org/10.1002/hed.21700>
 135. Silander E, Jacobsson I, Bertéus-Forslund H, Hammerlid E. Energy intake and sources of nutritional support in patients with head and neck cancer--a randomised longitudinal study. *Eur J Clin Nutr*. 2013;67:47-52.
<https://doi.org/10.1038/ejcn.2012.172>
 136. Axelsson L, Silander E, Nyman J, Bove M, Johansson L, Hammerlid E. Effect of prophylactic percutaneous endoscopic gastrostomy tube on swallowing in advanced head and neck cancer: a randomized controlled study. *Head Neck*. 2017;39:908-15.
<https://doi.org/10.1002/hed.24707>
 137. Brown TE, Banks MD, Hughes BGM, Lin CY, Kenny LM, Bauer JD. Randomised controlled trial of early prophylactic feeding vs standard care in patients with head and neck cancer. *Br J Cancer*. 2017;117:15-24.
<https://doi.org/10.1038/bjc.2017.138>
 138. Park RH, Allison MC, Lang J, Spence E, Morris AJ, Danesh BJ, et al. Randomised comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding in patients with persisting neurological dysphagia. *BMJ*. 1992;304:1406-9.
<https://doi.org/10.1136/bmj.304.6839.1406>
 139. Norton B, Homer-Ward M, Donnelly MT, Long RG, Holmes GK. A randomised prospective comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding after acute dysphagic stroke. *BMJ*. 1996;312:13-6.
<https://doi.org/10.1136/bmj.312.7022.13>
 140. Dennis M, Lewis S, Cranswick G, Forbes J. FOOD: a multicentre randomised trial evaluating feeding policies in patients admitted to hospital with a recent stroke. *Health Technol Assess* 2006;10:iii-iv, ix-x, 1-120.
<https://doi.org/10.3310/hta10020>
 141. Hamidon BB, Abdullah SA, Zawawi MF, Sukumar N, Aminuddin A, Raymond AA. A prospective comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding in patients with acute dysphagic stroke. *Med J Malaysia*. 2006;61:59-66.
 142. Corry J, Poon W, McPhee N, Milner AD, Cruickshank D, Porceddu SV, et al. Prospective study of percutaneous endoscopic gastrostomy tubes versus nasogastric tubes for enteral feeding in patients with head and neck cancer undergoing (chemo)radiation. *Head Neck*. 2009;31:867-76.
<https://doi.org/10.1002/hed.21044>
 143. Ambrosi D, Lee YT. Rehabilitation of swallowing disorders. In: Cifu DX, editors. *Braddom's physical medicine and rehabilitation*. Philadelphia: Elsevier; 2021. p. 53-67.e2.
 144. Murray J, Doeltgen S, Miller M, Scholten I. Does a water protocol improve the hydration and health status of individuals with thin liquid aspiration following stroke? A randomized controlled trial. *Dysphagia*. 2016;31:424-33.
<https://doi.org/10.1007/s00455-016-9694-x>
 145. Diniz PB, Vanin G, Xavier R, Parente MA. Reduced incidence of aspiration with spoon-thick consistency in stroke patients. *Nutr Clin Pract*. 2009;24:414-8.
<https://doi.org/10.1177/0884533608329440>
 146. Kyodo R, Kudo T, Horiuchi A, Sakamoto T, Shimizu T. Pureed diets containing a gelling agent to reduce the risk of aspiration in elderly patients with moderate to severe dysphagia: a randomized, crossover trial. *Medicine (Baltimore)*. 2020;99:e21165.
<https://doi.org/10.1097/md.00000000000021165>
 147. Robbins J, Gensler G, Hind J, Logemann JA, Lindblad AS, Brandt D, et al. Comparison of 2 interventions for liquid aspiration on pneumonia incidence: a randomized trial. *Ann Intern Med*. 2008;148:509-18. Erratum in: *Ann Intern Med*. 2008;148:715.
<https://doi.org/10.7326/0003-4819-148-7-200804010-00007>
 148. Teggart K, Phillips S, Ganann R, Moore C, Sihota D,

- Neil-Sztramko S, et al. Group-based nutrition interventions for community-dwelling older adults: a systematic review to inform community co-design. *Ann Fam Med*. 2022;20(Suppl 1):2875.
<https://doi.org/10.1370/afm.20.s1.2875>
149. Shimazu S, Yoshimura Y, Kudo M, Nagano F, Bise T, Shiraishi A, et al. Frequent and personalized nutritional support leads to improved nutritional status, activities of daily living, and dysphagia after stroke. *Nutrition*. 2021;83:111091.
<https://doi.org/10.1016/j.nut.2020.111091>
 150. Tappenden KA, Quatrara B, Parkhurst ML, Malone AM, Fanjiang G, Ziegler TR. Critical role of nutrition in improving quality of care: an interdisciplinary call to action to address adult hospital malnutrition. *J Acad Nutr Diet*. 2013;113:1219-37.
<https://doi.org/10.1016/j.jand.2013.05.015>
 151. Germain I, Dufresne T, Gray-Donald K. A novel dysphagia diet improves the nutrient intake of institutionalized elders. *J Am Diet Assoc*. 2006;106:1614-23.
<https://doi.org/10.1016/j.jada.2006.07.008>
 152. Reyes-Torres CA, Castillo-Martínez L, Reyes-Guerrero R, Ramos-Vázquez AG, Zavala-Solares M, Cassis-Nosthas L, et al. Design and implementation of modified-texture diet in older adults with oropharyngeal dysphagia: a randomized controlled trial. *Eur J Clin Nutr*. 2019;73:989-96.
<https://doi.org/10.1038/s41430-019-0389-x>
 153. Taylor KA, Barr SI. Provision of small, frequent meals does not improve energy intake of elderly residents with dysphagia who live in an extended-care facility. *J Am Diet Assoc*. 2006;106:1115-8.
 154. Shimizu A, Fujishima I, Maeda K, Wakabayashi H, Nishioka S, Ohno T, et al. Nutritional management enhances the recovery of swallowing ability in older patients with sarcopenic dysphagia. *Nutrients*. 2021;13:596.
 155. Zanini M, Bagnasco A, Catania G, Aleo G, Sartini M, Cristina ML, et al. A DEDICATED NUTRITIONAL CARE PROGRAM (NUTRICARE) to reduce malnutrition in institutionalised dysphagic older people: a quasi-experimental study. *J Clin Nurs*. 2017;26:4446-55.
 156. Wright L, Cotter D, Hickson M. The effectiveness of targeted feeding assistance to improve the nutritional intake of elderly dysphagic patients in hospital. *J Hum Nutr Diet*. 2008;21:555-62; quiz 564-5.
 157. Martens L, Cameron T, Simonsen M. Effects of a multidisciplinary management program on neurologically impaired patients with dysphagia. *Dysphagia*. 1990;5:147-51.
 158. An TJ, Myong JP, Lee YH, Kwon SO, Shim EK, Shin JH, et al. Continuing quality assessment program improves clinical outcomes of hospitalized community-acquired pneumonia: a nationwide cross-sectional study in Korea. *J Korean Med Sci*. 2022;37:e234.
<https://doi.org/10.3346/jkms.2022.37.e234>
 159. Hayashi M, Iwasaki T, Yamazaki Y, Takayasu H, Tateno H, Tazawa S, et al. Clinical features and outcomes of aspiration pneumonia compared with non-aspiration pneumonia: a retrospective cohort study. *J Infect Chemother*. 2014;20:436-42.
<https://doi.org/10.1016/j.jiac.2014.04.002>
 160. Feng MC, Lin YC, Chang YH, Chen CH, Chiang HC, Huang LC, et al. The mortality and the risk of aspiration pneumonia related with dysphagia in stroke patients. *J Stroke Cerebrovasc Dis*. 2019;28:1381-7.
<https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.02.011>
 161. Jeon I, Jung GP, Seo HG, Ryu JS, Han TR, Oh BM. Proportion of aspiration pneumonia cases among patients with community-acquired pneumonia: a single-center study in Korea. *Ann Rehabil Med*. 2019;43:121-8.
<https://doi.org/10.5535/arm.2019.43.2.121>
 162. Reza Shariatzadeh M, Huang JQ, Marrie TJ. Differences in the features of aspiration pneumonia according to site of acquisition: community or continuing care facility. *J Am Geriatr Soc*. 2006;54:296-302.
<https://doi.org/10.1111/j.1532-5415.2005.00608.x>
 163. van der Maarel-Wierink CD, Vanobbergen JN, Bronkhorst EM, Schols JM, de Baat C. Meta-analysis of dysphagia and aspiration pneumonia in frail elders. *J Dent Res*. 2011;90:1398-404.
<https://doi.org/10.1177/0022034511422909>
 164. Lo WL, Leu HB, Yang MC, Wang DH, Hsu ML. Dysphagia and risk of aspiration pneumonia: a non-randomized, pair-matched cohort study. *J Dent Sci*. 2019;14:241-7.
<https://doi.org/10.1016/j.jds.2019.01.005>
 165. Manabe T, Teramoto S, Tamiya N, Okochi J, Hizawa N. Risk factors for aspiration pneumonia in older adults. *PLoS One*. 2015;10:e0140060.
<https://doi.org/10.1371/journal.pone.0140060>
 166. Hunter KU, Lee OE, Lyden TH, Haxer MJ, Feng FY, Schipper M, et al. Aspiration pneumonia after chemo-intensity-modulated radiation therapy of oropharyngeal carcinoma and its clinical and dysphagia-related predictors. *Head Neck*. 2014;36:120-5.
<https://doi.org/10.1002/hed.23275>
 167. Watila MM, Nyandaiti YW, Balarabe SA, Bakki B, Alkali NH, Ibrahim A, et al. Aspiration pneumonia in patients with stroke in Northeast Nigeria. *Int J Stroke*. 2013;8:E16.
<https://doi.org/10.1111/ijis.12095>
 168. Xu Z, Gu Y, Li J, Wang C, Wang R, Huang Y, et al. Dysphagia and aspiration pneumonia in elderly hospitalization stroke patients: risk factors, cerebral infarction area comparison. *J Back Musculoskelet Rehabil*. 2019;32:85-91.
<https://doi.org/10.3233/bmr-170801>
 169. Sumi Y. [The significance of oral care in dysphagia patients]. *Nihon Ronen Igakkai Zasshi*. 2013;50:465-8. Japanese.
<https://doi.org/10.1152/ajpcell.00093.2008>
 170. Toniolo L, Cancellara P, Maccatrozzo L, Patruno M,

- Mascarello F, Reggiani C. Masticatory myosin unveiled: first determination of contractile parameters of muscle fibers from carnivore jaw muscles. *Am J Physiol Cell Physiol*. 2008;295:C1535-42.
https://doi.org/10.1152/ajpcell.00093.2008
171. Chen HJ, Chen JL, Chen CY, Lee M, Chang WH, Huang TT. Effect of an oral health programme on oral health, oral intake, and nutrition in patients with stroke and dysphagia in taiwan: a randomised controlled trial. *Int J Environ Res Public Health*. 2019;16:2228.
https://doi.org/10.3390/ijerph16122228
 172. Chippis E, Gatens C, Genter L, Musto M, Dubis-Bohn A, Gliemmo M, et al. Pilot study of an oral care protocol on poststroke survivors. *Rehabil Nurs*. 2014;39:294-304.
https://doi.org/10.1002/rnj.154
 173. Bai AV, Agostini F, Bernetti A, Mangone M, Fidenzi G, D'Urzo R, et al. State of the evidence about re-habilitation interventions in patients with dysphagia. *Eur J Phys Rehabil Med*. 2021;57:900-11.
https://doi.org/10.23736/s1973-9087.21.06716-2
 174. Kelley RE, Borazanci AP. Stroke rehabilitation. *Neurol Res*. 2009;31:832-40.
https://doi.org/10.1179/016164109x12445505689689
 175. Kristensen MB, Isenring E, Brown B. Nutrition and swallowing therapy strategies for patients with head and neck cancer. *Nutrition*. 2020;69:110548.
https://doi.org/10.1016/j.nut.2019.06.028
 176. Zheng L, Li Y, Liu Y. The individualized rehabilitation interventions for dysphagia: a multidisciplinary case control study of acute stroke patients. *Int J Clin Exp Med*. 2014;7:3789-94.
 177. Aoki S, Hosomi N, Hirayama J, Nakamori M, Yoshikawa M, Nezu T, et al. The multidisciplinary swallowing team approach decreases pneumonia onset in acute stroke patients. *PLoS One*. 2016;11:e0154608.
https://doi.org/10.1371/journal.pone.0154608
 178. Gandolfi M, Smania N, Bisoffi G, Squaquara T, Zuccher P, Mazzucco S. Improving post-stroke dysphagia outcomes through a standardized and multidisciplinary protocol: an exploratory cohort study. *Dysphagia*. 2014;29:704-12.
https://doi.org/10.1007/s00455-014-9565-2
 179. Starmer HM, Ayoub N, Byward C, Kizner J, Le Q, Hara W, et al. The impact of developing a speech and swallow rehab program: improving patient satisfaction and multidisciplinary care. *Laryngoscope*. 2017;127:2578-81.
https://doi.org/10.1002/lary.26695
- Hee Seung Yang, M.D., Ph.D.⁹,
Wonjae Cha, M.D., Ph.D.¹⁰, Ji Won Kim, M.D., Ph.D.¹¹, Byung-Mo Oh, M.D., Ph.D.¹²,
Han Gil Seo, M.D., Ph.D.¹², Min-Wook Kim, M.D., Ph.D.¹³, Hee-Soon Woo, O.T., Ph.D.¹⁴,
Sung-Jong Park, S.T.¹⁵, Sungju Jee, M.D.¹⁶,
Ju Sun Oh, M.D.¹⁷, Ki Deok Park, M.D., Ph.D.¹⁸,
Young Ju Jin, M.D.¹⁹, Sungjun Han, M.D.²⁰,
DooHan Yoo, O.T., Ph.D.²¹, Bo Hae Kim, M.D., Ph.D.²², Hyun Haeng Lee, M.D., M.S.²³,
Yeo Hyung Kim, M.D., Ph.D.²⁴,
Min-Gu Kang, M.D.²⁵, Eun-Jae Chung, M.D., Ph.D.²⁶, Bo Ryun Kim, M.D., Ph.D.²⁷,
Tae-Woo Kim, M.D.^{28,29}, Eun Jae Ko, M.D., Ph.D.⁸, Young Min Park, M.D., Ph.D.³⁰,
Hanaro Park, M.D.³¹, Min-Su Kim, M.D., Ph.D.³², Jungirl Seok, M.D.³³, Sun Im, M.D., Ph.D.³⁴, Sung-Hwa Ko, M.D., Ph.D.³⁵,
Seong Hoon Lim, M.D., Ph.D.³⁶, Kee Wook Jung, M.D., Ph.D.³⁷, Tae Hee Lee, M.D., Ph.D.³⁸, Bo Young Hong, M.D., Ph.D.³⁶,
Woojeong Kim, M.S., R.D.³⁹, Weon-Sun Shin, Ph.D.⁴⁰, Young Chan Lee, M.D.⁴¹,
Sung Joon Park, M.D.⁴², Jeonghyun Lim, Ph.D., R.D.⁴³, Youngkook Kim, M.D., Ph.D.⁴⁴,
Jung Hwan Lee, M.D., Ph.D.⁴⁵, Kang-Min Ahn, D.D.S., M.S.D., Ph.D.⁴⁶, Jun-Young Paeng, D.D.S., Ph.D.⁴⁷, JeongYun Park, R.N., Ph.D.⁴⁸,
Young Ae Song, M.S.N., R.N.⁴⁹,
Kyung Cheon Seo, M.D.⁵⁰, Chang Hwan Ryu, M.D.⁵¹, Jae-Keun Cho, M.D., Ph.D.⁵²,
Jee-Ho Lee, D.D.S.⁵³, Kyoung Hyo Choi, M.D., Ph.D.⁸

¹Department of Rehabilitation Medicine, Ewha Womans University Seoul Hospital, Ewha Womans University College of Medicine, Seoul, ²Department of Physical Medicine and Rehabilitation, Dongguk University Ilsan Hospital, Goyang, ³Department of Rehabilitation Medicine, CHA Bundang Medical Center, CHA University School of Medicine, Seongnam, ⁴Department of Otorhinolaryngology-Head and Neck Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, ⁵Department of Occupational Therapy, Asan Medical Center, University of Ulsan College of Medicine, Seoul, ⁶Department Audiology and Speech-Language Pathology, Daegu Catholic University, Gyeongsan, ⁷Department of Rehabilitation Medicine, International St. Mary's Hospital, Catholic Kwandong University College of Medicine, Incheon, ⁸Department of Rehabilitation Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, ⁹Department of Physical Medicine and Rehabilitation, Veterans Health Service Medical Center, Seoul, ¹⁰Department of Otorhinolaryngology-Head and Neck Surgery, Seoul National University Bundang Hospital, Seoul National University College of Medicine, Seongnam,

Seoyon Yang, M.D., Ph.D.¹, Jin-Woo Park, M.D., Ph.D.², Kyunghoon Min, M.D., Ph.D.³, Yoon Se Lee, M.D., Ph.D.⁴, Young-Jin Song, O.T.⁵, Seong Hee Choi, Ph.D.⁶, Doo Young Kim, M.D.⁷, Seung Hak Lee, M.D., Ph.D.⁸,

¹¹Department of Otolaryngology, Inha University College of Medicine, Incheon, ¹²Department of Rehabilitation Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, ¹³Department of Rehabilitation Medicine, Incheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Incheon, ¹⁴Department of Occupational Therapy, School of Medicine, Wonkwang University, Iksan, ¹⁵Department of Speech Therapy, Asan Medical Center, University of Ulsan College of Medicine, Seoul, ¹⁶Department of Rehabilitation Medicine, Chungnam National University Hospital, College of Medicine, Chungnam National University, Daejeon, ¹⁷Department of Rehabilitation Medicine, Seoul Medical Center, Seoul, ¹⁸Department of Rehabilitation Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon, ¹⁹Department of Otorhinolaryngology-Head and Neck Surgery, Kangwon National University Hospital, Kangwon National University College of Medicine, Chuncheon, ²⁰Department of Otolaryngology-Head and Neck Surgery, St. Vincent's Hospital, College of Medicine, The Catholic University of Korea, Seoul, ²¹Department of Occupational Therapy, Konyang University, Daejeon, ²²Department of Otorhinolaryngology-Head and Neck Surgery, Dongguk University Ilsan Hospital, Goyang, ²³Department of Rehabilitation Medicine, Konkuk University Medical Center, Seoul, ²⁴Department of Rehabilitation Medicine, College of Medicine, The Catholic University of Korea, Seoul, ²⁵Department of Physical Medicine and Rehabilitation, Dong-A University College of Medicine, Busan, ²⁶Department of Otorhinolaryngology-Head and Neck Surgery, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, ²⁷Department of Physical Medicine and Rehabilitation, Korea University Anam Hospital, Korea University College of Medicine, Seoul, ²⁸Department of Rehabilitation Medicine, National Traffic Injury Rehabilitation Hospital, Yangpyeong, ²⁹Department of Rehabilitation Medicine, Seoul National University College of Medicine, Seoul, ³⁰Department of Otorhinolaryngology, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, ³¹Department of Otorhinolaryngology-Head and Neck Surgery, Samsung Changwon Hospital, Sungkyunkwan University School of Medicine, Changwon, ³²Department of Otorhinolaryngology-Head and Neck Surgery, CHA Bundang

Medical Center, CHA University School of Medicine, Seongnam, ³³Department of Otorhinolaryngology-Head and Neck Surgery, Seoul National University Hospital, Seoul, ³⁴Department of Rehabilitation Medicine, Bucheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Bucheon, ³⁵Department of Rehabilitation Medicine, Pusan National University & Pusan National University Yangsan Hospital, Yangsan, ³⁶Department of Rehabilitation Medicine, St. Vincent's Hospital, College of Medicine, The Catholic University of Korea, Seoul, ³⁷Department of Gastroenterology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, ³⁸Department of Gastroenterology, Konyang University College of Medicine, Daejeon, ³⁹Department of Nutrition & Food Control, Gangnam Severance Hospital, Seoul, ⁴⁰Department of Food & Nutrition, College of Human Ecology, Hanyang University, Seoul, ⁴¹Department of Otorhinolaryngology-Head and Neck Surgery, Kyung Hee University Hospital at Gangdong, Kyung Hee University School of Medicine, Seoul, ⁴²Department of Otorhinolaryngology-Head and Neck Surgery, Chung-Ang University Gwangmyeong Hospital, Chung-Ang University College of Medicine, Gwangmyeong, ⁴³Department of Food Service & Nutrition Care, Seoul National University Hospital, Seoul, ⁴⁴Department of Rehabilitation Medicine, Yeouido St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, ⁴⁵Namdarun Rehabilitation Clinic, Yongin, ⁴⁶Department of Oral and Maxillofacial Surgery, University of Ulsan College of Medicine, Seoul, ⁴⁷Department of Oral and Maxillofacial Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, ⁴⁸Department of Clinical Nursing, University of Ulsan, Seoul, ⁴⁹Department of Nursing, Seoul National University Bundang Hospital, Seongnam, ⁵⁰Department of Physical Medicine and Rehabilitation, Korea University Anam Hospital, Seoul, ⁵¹Department of Otolaryngology-Head and Neck Surgery, Center for Thyroid Cancer, Research Institute and Hospital, National Cancer Center, Goyang, ⁵²Department of Otorhinolaryngology-Head and Neck Surgery, Inje University Ilsan Paik Hospital, Inje University College of Medicine, Goyang, ⁵³Department of Oral and Maxillofacial Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea